

International Textbook of Diabetes Mellitus, 4th Ed., Chapter 4: Development and Maintenance of the Islet β Cell

References:

1. Andrew A: An experimental investigation into the possible neural crest origin of pancreatic APUD (islet) cells. *Journal of Embryology and Experimental Morphology* 1976;35:577–593.
2. Schier AF, Shen MM: Nodal signalling in vertebrate development. *Nature* 2000;403:385–389.
3. Behringer RR, Bradley A, Liu P, et al.: Requirement for Wnt3 invertebrate axis formation. *Nature Genetics* 1999;22:361–365.
4. Xanthos JB, Kofron M, Wylie C, Heasman J: Maternal VegT is the initiator of a molecular network specifying endoderm in *Xenopus laevis*. *Development* 2001;128:167–180.
5. Schier AF, Talbot WS: Molecular genetics of axis formation in zebrafish. *Annual Review of Genetics* 2005;39:561–613.
6. Grapin-Botton A, Constam D: Evolution of the mechanisms and molecular control of endoderm formation. *Mechanisms of Development* 2007;124:253–278.
7. Stainier DYR: A glimpse into the molecular entrails of endoderm formation. *Genes & Development* 2002;16(8):893–907.
8. Zorn AM, Wells JM: Vertebrate endoderm development and organ formation. *Annual Review of Cell and Developmental Biology* 2009;25:221–251.
9. Martinez Barbera JP, Clements M, Thomas P, et al.: The homeobox gene *Hex* is required in definitive endodermal tissues for normal forebrain, liver and thyroid formation. *Development* 2000;127:2433–2445.
10. Que J, Okubo T, Goldenring JR, et al.: Multiple dose-dependent roles for *Sox2* in the patterning and differentiation of anterior foregut endoderm. *Development* 2007;134:2521–2531.
11. Zaret KS: Genetic programming of liver and pancreas progenitors: lessons for stem-cell differentiation. *Nature Reviews Genetics* 2008;9:329–340.
12. Zaret KS, Grompe M: Generation and regeneration of cells of the liver and pancreas. *Science* 2008;322:1490–1494.
13. Shin D, Shin CH, Tucker J, et al.: Bmp and Fgf signaling are essential for liver specification in zebrafish. *Development* 2007; 134:2041–2050.
14. Molotkov A, Molotkova N, Duester G: Retinoic acid generated by *Raldh2* in mesoderm is required for mouse dorsal endodermal pancreas development. *Developmental Dynamics* 2005; 232:950–957.
15. Stafford D, White RJ, Kinkel MD, et al.: Retinoids signal directly to zebrafish endoderm to specify insulin-expressing β -cells. *Development* 2006;133:949–956.
16. Burlison JS, Long Q, Fujitani Y, et al.: *Pdx-1* and *Ptf1a* concurrently determine fate specification of pancreatic multipotent progenitor cells. *Developmental Biology* 2008;316:74–86.
17. Hebrok M, Kim SK, Melton DA: Notochord repression of endodermal Sonic hedgehog permits pancreas development. *Genes & Development* 1998;12:1705–1713.
18. Lau J, Hebrok M: Hedgehog signaling in pancreas epithelium regulates embryonic organ formation and adult β -cell function. *Diabetes* 2010;59:1211–1221.
19. Stafford D, Prince VE: Retinoic acid signaling is required for a critical early step

- in zebrafish pancreatic development. *Current Biology* 2002;12:1215–1220.
20. Chung W-S, Stainier DYR: Intra-endodermal interactions are required for pancreatic β cell induction. *Developmental Cell* 2008; 14:582–593.
 21. Field HA, Dong PDS, Beis D, Stainier DYR: Formation of the digestive system in zebrafish. ii. pancreas morphogenesis. *Developmental Biology* 2003;261:197–208.
 22. Gittes GK, Galante PE, Hanahan D, et al.: Lineage-specific morphogenesis in the developing pancreas: role of mesenchymal factors. *Development* 1996;122:439–447.
 23. Landsman L, Nijagal A, Whitchurch TJ, et al.: Pancreatic mesenchyme regulates epithelial organogenesis throughout development. *PLoS Biology* 2011;9:e1001143.
 24. Bhushan A, Itoh N, Kato S, et al.: Fgf10 is essential for maintaining the proliferative capacity of epithelial progenitor cells during early pancreatic organogenesis. *Development* 2001;128:5109–5117.
 25. Norgaard GA, Jensen JN, Jensen J: FGF10 signaling maintains the pancreatic progenitor cell state revealing a novel role of Notch in organ development. *Developmental Biology* 2003;264:323–338.
 26. Villasenor A, Chong DC, Henkemeyer M, Cleaver O: Epithelial dynamics of pancreatic branching morphogenesis. *Development* 2010;137:4295–4305.
 27. Magenheim J, Ilovich O, Lazarus A, et al.: Blood vessels restrain pancreas branching, differentiation and growth. *Development* 2011;138:4743–4752.
 28. Stanger BZ, Tanaka AJ, Melton DA: Organ size is limited by the number of embryonic progenitor cells in the pancreas but not the liver. *Nature* 2007;445:886–891.
 29. Cano DA, Hebrok M, Zenker M: Pancreatic development and disease. *Gastroenterology* 2007;132:745–762.
 30. Pictet R, Rutter WJ: Development of the embryonic endocrine pancreas. In: *Handbook of Physiology*. Baltimore, MD: Williams and Wilkins, 1972, pp. 25–66.
 31. Zhou Q, Law AC, Rajagopal J, et al.: A multipotent progenitor domain guides pancreatic organogenesis. *Developmental Cell* 2007;13:103–114.
 32. Jorgensen MC, Ahnfelt-Ronne J, Hald J, et al.: An illustrated review of early pancreas development in the mouse. *Endocrine Reviews* 2007; 28:685–705.
 33. Schaffer AE, Freude KK, Nelson SB, Sander M: Nkx6 transcription factors and Ptf1a function as antagonistic lineage determinants in multipotent pancreatic progenitors. *Developmental Cell* 2010; 18:1022–1029.
 34. Wang S, Yan J, Anderson DA, et al.: Neurog3 gene dosage regulates allocation of endocrine and exocrine cell fates in the developing mouse pancreas. *Developmental Biology* 2010;339:26–37.
 35. Gradwohl G, Dierich A, LeMeur M, Guillemot F: Neurogenin3 is required for the development of the four endocrine cell lineages of the pancreas. *PNAS* 2000;97:1607–1611.
 36. Gu G, Dubauskaite J, Melton DA: Direct evidence for the pancreatic lineage: NGN3+ cells are islet progenitors and are distinct from duct progenitors. *Development* 2002;129:2447–2457.
 37. Heller RS, Jenny M, Collombat P, et al.: Genetic determinants of pancreatic ϵ -cell development. *Developmental Biology* 2005;286:217–224.
 39. Gouzi M, Kim YH, Katsumoto K, et al.: Neurogenin3 initiates stepwise delamination of differentiating endocrine cells during pancreas development. *Developmental Dynamics* 2011;240:589–604.

40. Johansson KA, Dursun U, Jordan N, et al.: Temporal control of neurogenin3 activity in pancreas progenitors reveals competence windows for the generation of different endocrine cell types. *Developmental Cell* 2007;12:457–465.
41. Desgraz R, Herrera PL: Pancreatic neurogenin 3-expressing cells are unipotent islet precursors. *Development* 2009;136:3567–3574.
42. Wang X, Chu LT, He J, et al.: A novel zebrafish bHLH gene, neurogenin3, is expressed in the hypothalamus. *Gene* 2001;275:47–55.
43. Pinney SE, Oliver-Krasinski J, Ernst L, et al.: Neonatal diabetes and congenital malabsorptive diarrhea attributable to a novel mutation in the human neurogenin-3 gene coding sequence. *Journal of Clinical Endocrinology & Metabolism* 2011;96:1960–1965.
44. Wang J, Cortina G, Wu SV, et al.: Mutant neurogenin-3 in congenital malabsorptive diarrhea. *New England Journal of Medicine* 2006;355:270–280.
45. St-Onge L, Sosa-Pineda B, Chowdhury K, et al.: Pax6 is required for differentiation of glucagon-producing alpha-cells in mouse pancreas. *Nature* 1997;387:406–409.
46. Shimomura H, Sanke T, Hanabusa T, et al.: Nonsense mutation of islet-1 gene (Q310X) found in a type 2 diabetic patient with a strong family history. *Diabetes* 2000;49:1597–1600.
47. Naya FJ, Huang HP, Qiu Y, et al.: Diabetes, defective pancreatic morphogenesis, and abnormal enteroendocrine differentiation in BETA2/neuroD-deficient mice. *Genes & Development* 1997; 11:2323–2334.
48. Artner I, Bianchi B, Raum JC, et al.: MafB is required for islet beta cell maturation. *PNAS* 2007;104:3853–3858.
49. Zhang C, Moriguchi T, Kajihara M, et al.: MafA is a key regulator of glucose-stimulated insulin secretion. *Molecular and Cellular Biology* 2005;25:4969–4976.
50. Jonsson J, Carlsson L, Edlund T, Edlund H: Insulin-promoter factor 1 is required for pancreas development in mice. *Nature* 1994;371:606–609.
52. Brissova M, Shiota M, Nicholson WE, et al.: Reduction in pancreatic transcription factor PDX-1 impairs glucose-stimulated insulin secretion. *The Journal of Biological Chemistry* 2002;277:11225–11232.
53. Babu DA, Deering TG, Mirmira RG: A feat of metabolic proportions: Pdx1 orchestrates islet development and function in the maintenance of glucose homeostasis. *Molecular Genetics and Metabolism* 2007;92:43–55.
54. Mastracci TL, Wilcox CL, Arnes L, et al.: Nkx2.2 and Arx genetically interact to regulate pancreatic endocrine cell development and endocrine hormone expression. *Developmental Biology* 2011;359:1–11.
55. Prado CL, Pugh-Bernard AE, Elghazi L, et al.: Ghrelin cells replace insulin-producing beta cells in two mouse models of pancreas development. *PNAS* 2004;101:2924–2929.
56. Fajans SS, Bell GI, Polonsky KS: Molecular mechanisms and clinical pathophysiology of maturity-onset diabetes of the young. *New England Journal of Medicine* 2001;345:971–980.
57. LangoAllen H, Flanagan SE, Shaw-Smith C, et al.: GATA6 haploinsufficiency causes pancreatic agenesis in humans. *Nature Genetics* 2011;44:20–22.
58. Pearl EJ, Jarikji Z, Horb ME: Functional analysis of Rfx6 and mutant variants associated with neonatal diabetes. *Developmental Biology* 2011;351:135–145.
59. Sellick GS, Barker KT, Stolte-Dijkstra I, et al.: Mutations in PTF1A cause pancreatic and cerebellar agenesis. *Nature Genetics* 2004;36:1301–1305.
60. Senee V, Chelala C, Duchatelet S, et al.: Mutations in GLIS3 are responsible for a rare syndrome with neonatal diabetes mellitus and congenital hypothyroidism.

- Nature Genetics 2006;38:682–687.
61. Zhou Q, Brown J, Kanarek A, et al.: In vivo reprogramming of adult pancreatic exocrine cells to β -cells. *Nature* 2008;455:627–632.
 62. Thorel F, Népoté V, Avril I, et al.: Conversion of adult pancreatic alpha-cells to beta-cells after extreme β -cell loss. *Nature* 2010;464:1149–1154.
 63. 2010;464:1149–1154.
 64. Ogihara T, Mirmira RG: An islet in distress: β cell failure in type 2 diabetes. *Journal of Diabetes Investigation* 2010;1:123–133.
 65. Bonner-Weir S: beta-Cell turnover: its assessment and implications. *Diabetes* 2001;50 (Suppl 1):S20–24.
 66. Scaglia L, Cahill CJ, Finegood DT, Bonner-Weir S: Apoptosis participates in the remodeling of the endocrine pancreas in the neonatal rat. *Endocrinology* 1997;138:1736–1741.
 67. Teta M, Long SY, Wartschow LM, et al.: Very slow turnover of beta-cells in aged adult mice. *Diabetes* 2005;54:2557–2567.
 68. Gunasekaran U, Hudgens CW, Wright BT, et al.: Differential regulation of embryonic and adult β cell replication. *Cell Cycle* 2012;11:2431–2442.
 69. Meier JJ, Butler AE, Saisho Y, et al.: Beta-cell replication is the primary mechanism subserving the postnatal expansion of β -cell mass in humans. *Diabetes* 2008;57:1584–1594.
 70. Withers DJ, Gutierrez JS, Towery H, et al.: Disruption of IRS-2 causes type 2 diabetes in mice. *Nature* 1998;391:900–904.
 71. Assmann A, Hinault C, Kulkarni RN: Growth factor control of pancreatic islet regeneration and function. *Pediatric Diabetes* 2009; 10:14–32.
 72. Dhawan S, Georgia S, Bhushan A: Formation and regeneration of the endocrine pancreas. *Current Opinion in Cell Biology* 2007; 19:634–645.
 73. Mathis D, Vence L, Benoist C: β -Cell death during progression to diabetes. *Nature* 2001;414:792–798.
 74. Carrington EM, Kos C, Zhan Y, et al.: Reducing or increasing β -cell apoptosis without inflammation does not affect diabetes initiation in neonatal NOD mice. *European Journal of Immunology* 2011;41:2238–2247.
 75. Bonner-Weir S: Perspective: postnatal pancreatic beta cell growth. *Endocrinology* 2000;141:1926–1929.
 76. Montanya E, Nacher V, Biarnés M, Soler J: Linear correlation between beta-cell mass and body weight throughout the lifespan in Lewis rats: role of beta-cell hyperplasia and hypertrophy. *Diabetes* 2000;49:1341–1346.
 77. Perl S, Kushner JA, Buchholz BA, et al.: Significant human beta-cell turnover is limited to the first three decades of life as determined by in vivo thymidine analog incorporation and radiocarbon dating. *Journal of Clinical Endocrinology & Metabolism* 2010;95: E234–239.
 78. Ogilvie RF: The islands of Langerhans in 19 cases of obesity. *Journal of Pathology and Bacteriology* 1933;37:473–481.
 79. Sachdeva MM, Claiborn KC, Khoo C, et al.: Pdx1 (MODY4) regulates pancreatic beta cell susceptibility to ER stress. *PNAS* 2009; 106:19090–19095.
 80. Bonner-Weir S, Baxter LA, Schuppin GT, Smith FE: A second pathway for regeneration of adult exocrine and endocrine
 81. pancreas. A possible recapitulation of embryonic development. *Diabetes* 1993;42:1715–1720.
 82. Dor Y, Brown J, Martinez OI, Melton DA: Adult pancreatic beta-cells are formed by self-duplication rather than stem-cell
 83. differentiation. *Nature* 2004;429:41–46.

84. Teta M, Rankin MM, Long SY, et al.: Growth and regeneration of adult beta cells does not involve specialized progenitors. *Developmental Cell* 2007;12:817–826.
85. Smukler SR, Arntfield ME, Razavi R, et al.: The adult mouse and human pancreas contain rare multipotent stem cells that express insulin. *Cell Stem Cell* 2011;8:281–293.
86. Desai BM, Oliver-Krasinski J, De Leon DD, et al.: Preexisting pancreatic acinar cells contribute to acinar cell, but not islet
87. beta cell, regeneration. *Journal of Clinical Investigation* 2007;117: 971–977.
88. Gregg BE, Moore PC, Demozay D, et al.: Formation of a human β -cell population within pancreatic islets is set early in life. *Journal of Clinical Endocrinology & Metabolism* 2012;97:3197–3206.
89. Alonso LC, Yokoe T, Zhang P, et al.: Glucose infusion in mice: a new model to induce beta-cell replication. *Diabetes* 2007;56: 1792–1801.
90. Kushner JA, Ciemerych MA, Sicinska E, et al.: Cyclins D2 and D1 are essential for postnatal pancreatic beta-cell growth. *Molecular Cell Biology* 2005;25:3752–3762.
91. Rane SG, Dubus P, Mettus RV, et al.: Loss of Cdk4 expression causes insulin-deficient diabetes and Cdk4 activation results in β -islet cell hyperplasia. *Nature Genetics* 1999;22:44–52.
92. Fiaschi-Taesch N, Bigatel TA, Sicari B, et al.: Survey of the human pancreatic β -cellG1/S proteome reveals a potential therapeutic role for Cdk-6 and Cyclin D1 in enhancing human β -cell replication and function in vivo. *Diabetes* 2009;58:882–893.
93. Krishnamurthy J, Ramsey MR, Ligon KL, et al.: p16INK4a induces an age-dependent decline in islet regenerative potential. *Nature* 2006;443:453–457.
94. Marx SJ: Molecular genetics of multiple endocrine neoplasia types 1 and 2. *Nature Reviews Cancer* 2005;5:367–375.
95. Garcia-Ocana A, Alonso LC: Glucose mediated regulation of β cell proliferation. *Open Endocrinology Journal* 2010;4:55–65.
96. Swenne I: Pancreatic beta-cell growth and diabetes mellitus. *Diabetologia* 1992;35:193–201.
97. Wu Y, Liu C, Sun H, et al.: Growth hormone receptor regulates β cell hyperplasia and glucose-stimulated insulin secretion in obese mice. *Journal of Clinical Investigation* 2011;121:2422–2426.
98. Vasavada RC, González-Pertusa JA, Fujinaka Y, et al.: Growth factors and beta cell replication. *International Journal of Biochemistry & Cell Biology* 2006;38:931–950.
99. Pissios P, Ozcan U, Kokkotou E, et al.: Melanin concentrating hormone is a novel regulator of islet function and growth. *Diabetes* 2007;56:311–319.
100. Kim H, Toyofuku Y, Lynn FC, et al.: Serotonin regulates pancreatic β cell mass during pregnancy. *Nature Medicine* 2010;16:804–808.
101. Soltani N, Qiu H, Aleksic M, et al.: GABA exerts protective and regenerative effects on islet beta cells and reverses diabetes. *PNAS* 2011;108:11692–11697.
102. Islam MS, Sjöholm A, Emilsson V: Fetal pancreatic islets express functional leptin receptors and leptin stimulates proliferation of fetal islet cells. *International Journal of Obesity and Related Metabolic Disorders* 2000;24:1246–1253.
103. Wijesekara N, Krishnamurthy M, Bhattacharjee A, et al.: Adiponectin-induced ERK and Akt phosphorylation protects against pancreatic beta cell apoptosis and increases insulin gene expression and secretion. *The Journal of*

- Biological Chemistry 2010;285:33623–33631.
104. Baggio LL, Drucker DJ: Biology of incretins: GLP-1 and GIP. *Gastroenterology* 2007;132:2131–2157.
 105. Suarez-Pinzon WL, Lakey JRT, Brand SJ, Rabinovitch A: Combination therapy with epidermal growth factor and gastrin induces neogenesis of human islet {beta}-cells from pancreatic duct cells and an increase in functional β -cell mass. *Journal of Clinical Endocrinology & Metabolism* 2005;90:3401–3409.
 106. Lavine JA, Attie AD: Gastrointestinal hormones and the regulation of β -cell mass. *Annals of the N Y Academy of Sciences* 2010;1212:41–58.
 107. Hart AW, Baeza N, Apelqvist A, Edlund H: Attenuation of FGF signalling in mouse beta-cells leads to diabetes. *Nature* 2000; 408:864–868.
 108. Brissova M, Shostak A, Shiota M, et al.: Pancreatic islet production of vascular endothelial growth factor-A is essential for islet vascularization, revascularization, and function. *Diabetes* 2006;55:2974–2985.
 109. Clemens TL, Karsenty G: The osteoblast: an insulin target cell controlling glucose homeostasis. *Journal of Bone and Mineral Research* 2011;26:677–680.