

How To Get Prior Approvals For Drugs That Are Not Covered

Created and shared by Stanley Schwartz, MD, FACP. FACE

For Prior Authorization

- ☐ had hypoglycemia on SU/glynide
- ☐ had hypoglycemia on insulin - stopping insulin
- ☐ high risk if patient gets hypoglycemia
  - ☐ drives a lot
  - ☐ walks up and down stairs
  - ☐ fall risk: morbid obesity; ankle/knee/hip problem; DM neuropathy
- ☐ ASCVD, arrhythmia, angina, S/P, MI
- ☐ TIA, CVA, carotid bruit
- ☐ watches little children
- ☐ electric tools (including blow dryer, curling iron)
- ☐ uses guns or other dangerous profession
- ☐ uses electric/gas ranges

© 2018, Diabetes in Control

How to Use Prior Authorization Stickers

1. attach to script/ paper chart, create as macro for e-scripts

2. If true, I check off first 1 or 2 boxes ('stopping insulin' kept if not approved for insulin combo, but crossed out if med can be given with insulin.)

3. Check off third box for everyone, with appropriate checks in other boxes to support contention

NOTE: Pharmacists often call insurer to avoid need for 'prior auth' as you already justified use, or if get form, sect'y can easily fill-out/submit

ENJOY!

© 2018, Diabetes in Control