International Textbook of Diabetes Mellitus, 4th Ed., Chapter 3: Epidemiology and Geography of Type 2 Diabetes Mellitus

References:

1. Zimmet P, Alberti KG, Shaw J: Global and societal implications of the diabetes epidemic. Nature 2001;414(6865):782–787.
2. Joslin EP: The prevention of diabetes mellitus. JAMA 1921; 76:79–84.
3. King H, Aubert RE, Herman WH: Global burden of diabetes, 1995–2025: prevalence, numerical estimates, and projections. Diabetes Care 1998;21:1414–1431.
4. National Diabetes Data Group: Classification and diagnosis of diabetes mellitus and other categories of glucose intolerance. Diabetes 1979;28(12):1039–1057.
5. World Health Organization:WHO Expert Committee on Diabetes Mellitus: Second Report. Technical Report Series No. 646. Geneva: WHO, 1980.
6. World Health Organization: Diabetes Mellitus: Report of a Study Group. Technical Report Series No. 727. Geneva:WHO, 1985.
7. Expert Committee on the Diagnosis and Classification of Diabetes Mellitus: Report of the Expert Committee on the Diagnosis and Classification of Diabetes Mellitus. Diabetes Care 1997;20(7): 1183–1197.
8. WHO Consultation: Definition, Diagnosis and Classification of Diabetes Mellitus and Its Complications. Part 1: Diagnosis and Classification of Diabetes Mellitus. Report No. 99.2. Geneva: WHO, 1999.
9. The DECODE Study Group: Is the current definition for diabetes relevant to mortality risk from all-cause, cardiovascular and non-cardiovascular disease? Diabetes Care 2003;26(3):688–696.
10. The DECODE Study Group: Age- and sex-specific prevalences of diabetes and impaired glucose regulation in 13 European cohorts. Diabetes Care 2003;26(1):61–69.
11. Carstensen B, Lindström J, Sundvall J, Borch-Johnsen K, Tuomilehto J, and the DPS Study Group. Measurement of blood glucose: comparison between different types of specimens. Ann Clin Biochem 2008;45:140–148.
12. The DECODA Study Group: Age- and sex-specific prevalences of diabetes and impaired glucose regulation in 11 Asian cohorts. Diabetes Care 2003;26(6):1770–1780.
13. Rathmann W, Haastert B, Icks A, et al.: High prevalence of undiagnosed diabetes mellitus in Southern Germany: target populations for efficient screening. The KORA Survey 2000. Diabetologia 2003;46(2):182–189.
14. Satman I, Omer B, Tutuncu Y, et al. HYPERLINK "http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3604592/" Twelve-year trends in the prevalence and risk factors of diabetes and prediabetes in Turkish adults. Eur J Epidemiol 2013;28(2):169–180.
15. Satman I, Yilmaz T, Sengul A, et al.: Population-based study of diabetes and risk characteristics in Turkey: results of the Turkish Diabetes Epidemiology Study (TURDEP). Diabetes Care 2002;25(9):1551–1556.
16. Harris MI, Flegal KM, Cowie CC, et al.: Prevalence of diabetes, impaired fasting glucose, and impaired glucose tolerance in U.S. adults. Diabetes Care 1998;21:518–524.
17. King H, Rewers M, WHO Ad Hoc Diabetes Reporting Group: Global estimates for prevalence of diabetes mellitus and impaired glucose tolerance in adults. Diabetes Care 1993;16:157–177.
18. Grandinetti A, Chang HK, Mau MK, et al.: Prevalence of glucose intolerance among Native Hawaiians in two rural communities. Native Hawaiian Health Research (NHHR) Project. Diabetes Care 1998;21(4):549–554.
19. Dunstan DW, Zimmet PZ, Welborn TA, et al.: The rising prevalence of diabetes and impaired glucose tolerance: the Australian Diabetes, Obesity and Lifestyle Study. Diabetes Care 2002; 25(5):829–834.
20. Park Y, Lee H, Koh CS, et al.: Prevalence of diabetes and IGT in Yonchon County, South Korea. Diabetes Care 1995;18(4):545–548.

21. Shera AS, Rafique G, Khawaja IA, et al.: Pakistan National Diabetes Survey: prevalence of glucose intolerance and associated factors in Baluchistan province. Diabetes Res Clin Pract 1999;44 (1): 49–58.
22. Shera AS, Rafique G, Khwaja IA, et al.: Pakistan National Diabetes Survey: prevalence of glucose intolerance and associated factors in Shikarpur, Sindh Province. Diabet Med 1995;12(12):1116–1121.
23. Shera AS, Rafique G, Khwaja IA, et al.: Pakistan National Diabetes Survey: prevalence of glucose intolerance and associated factors in North West at Frontier Province (NWFP) of Pakistan. J Pak Med Assoc 1999;49(9):206–211.
24. King H,Djumaeva S,Abdullaev B,GacicDobo M: Epidemiology of glucose intolerance and associated factors in Uzbekistan: a survey in Sirdaria province. Diabetes Res Clin Pract 2002;55(1):19–27.
25. Suvd J, Gerel B, Otgooloi H, et al.: Glucose intolerance and associated factors in Mongolia: results of a national survey. Diabet Med 2002;19(6):502–508.
26. Pan XR, Yang WY, Li GW, Liu J: Prevalence of diabetes and its risk factors in China, 1994. Diabetes Care 1997;20:1664–1669.
27. Yang W, Lu J,Weng J, et al.: Prevalence of diabetes among men and women in China. N Engl J Med. 2010;362(12):1090–1101.
28. Zimmet P, King H, Taylor R, et al.: The high prevalence of diabetes mellitus impaired glucose tolerance and diabetic retinopathy in Nauru—the 1982 survey. Diabetes Res 1984;1(1):13–18.
29. Dowse GK, Zimmet PZ, Finch CF, Collins VR: Decline in incidence of epidemic glucose intolerance in Nauruans: implications for the "thrifty genotype." Am J Epidemiol 1991;133(11): 1093–1104.
30. King H, Finch C, Collins A, et al.: Glucose tolerance in Papua New Guinea: ethnic differences, association with environmental and behavioural factors and the possible emergence of glucose intolerance in a highland community. Med J Aust 1989;151(4):204–210.
31. Collins VR, Dowse GK, Toelupe PM, et al.: Increasing prevalence of NIDDM in the Pacific island population of Western Samoa over a 13-year period. Diabetes Care 1994;17(4):288–296.
32. Colagiuri S, Colagiuri R, Na'ati S, et al.: The prevalence of diabetes in the kingdom of Tonga. Diabetes Care 2002;25(8):1378–1383.
33. Al-Nuaim AR: Prevalence of glucose intolerance in urban and rural communities in Saudi Arabia. Diabet Med 1997;14(7):595–602.
34. Asfour MG, Lambourne A, Soliman A, et al.: High prevalence of diabetes mellitus and impaired glucose tolerance in the Sultanate of Oman: results of the 1991 National Survey. Diabet Med 1995;12(12):1122–1125.
35. Husseini A, Abdul-Rahim H, Awartani F, et al.: Prevalence of diabetes mellitus and impaired glucose tolerance in a rural Palestinian population. East Mediterr Health J 2000;6(5–6):1039–1045.
36. Abdella N, Al Nakhi A, Al Arouj M, et al.: Impact of the 1997 American Diabetes Association criteria on classification of glucose intolerance among Kuwaitis below 50 years of age. Acta Diabetol 1999;36(3):133–140.
37. Dowse GK, Gareeboo H, Zimmet PZ, et al.: High prevalence of NIDDM and impaired glucose tolerance in Indian, Creole, and Chinese Mauritians. Diabetes 1990;39(3):390–396.
38. Swai AB,McLarty DG, Sherrif F, et al.: Diabetes and impaired glucose tolerance in an Asian community in Tanzania. Diabetes Res Clin Pract 1990;8(3):227–234.
39. McLarty DG, Swai AB, Kitange HM, et al.: Prevalence of diabetes and impaired glucose tolerance in rural Tanzania. Lancet 1989;1(8643):871–875.
40. Santos JL, Perez-Bravo F, Carrasco E, et al.: Low prevalence of type 2 diabetes despite a high average body mass index in the Aymara natives from Chile. Nutrition 2001;17(4):305–309.
41. Perez-Bravo F, Carrasco E, Santos JL, et al.: Prevalence of type 2 diabetes and obesity in rural Mapuche population from Chile. Nutrition 2001;17(3):236–238.
42. Larenas G, Arias G, Espinoza O: Prevalence of diabetes mellitus in an indigenous (Mapuche) community of the IXregion in Chile. Rev Med Chile 1985;113:1121–1125.
43. Dogadin SA,Mashtakov BP, Taranushenko TE: Prevalence of type 2 diabetes in northern populations of Siberia. Int J Circumpolar Health 2001;60(2):205–210.

44. The DECODE Study Group:Glucose tolerance and mortality: comparison of WHO and American Diabetes Association diagnostic criteria. Lancet 1999;354(9179):617–621.
45. The DECODE Study Group: Glucose tolerance and cardiovascular mortality: comparison of fasting and 2-hour diagnostic criteria. Arch Intern Med 2001;161(3):397–405.
46. Qiao Q, Pyorala K, Pyorala M, et al.: Two-hour glucose is a better risk predictor for incident coronary heart disease and cardiovascular mortality than fasting glucose. Eur Heart J 2002;23(16):1267–1275.
47. Tominaga M, Eguchi H, Manaka H, et al.: Impaired glucose tolerance is a risk factor for cardiovascular disease, but not impaired fasting glucose: the Funagata Diabetes Study. Diabetes Care 1999;22(6):920–924.
48. Shaw JE, Zimmet PZ, deCourten M, et al.: Impaired fasting glucose or impaired glucose tolerance. What best predicts future diabetes in Mauritius? Diabetes Care 1999;22(3):399–402.
49. Vaccaro O, Ruffa G, Imperatore G, et al.: Risk of diabetes in the new diagnostic category of impaired fasting glucose: a prospective analysis. Diabetes Care 1999;22(9):1490–1493.
50. Gabir MM, Hanson RL, Dabelea D, et al.: The 1997 American Diabetes Association and 1999 World Health Organization criteria for hyperglycemia in the diagnosis and prediction of diabetes. Diabetes Care 2000;23(8):1108–1112.
51. de Vegt F, Dekker JM, Jager A, et al.: Relation of impaired fasting and postload glucose with incident type 2 diabetes in a Dutch population: the Hoorn Study. JAMA 2001;285(16):2109–2113.
52. Qiao Q, Lindströ m J, Valle TT, Tuomilehto J. Progression to clinically diagnosed and treated diabetes from impaired glucose tolerance and impaired fasting glycaemia. Diabet Med 2003; 20:1027–1033.
53. Unwin N, Shaw J, Zimmet P, Alberti KG: Impaired glucose tolerance and impaired fasting glycaemia: the current status on definition and intervention. Diabet Med 2002;19(9):708–723.
54. The DECODE Study Group on behalf of the European Diabetes Epidemiology Study Group: Will new diagnostic criteria for diabetes mellitus change phenotype of patients with diabetes? Reanalysis of European epidemiological data. BMJ 1998; 317(7155):371–375.
55. The DECODE Study Group: Is fasting glucose sufficient to define diabetes? Epidemiological data from 20 European studies. Diabetologia 1999;42(6):647–654.
56. Qiao Q, Nakagami T, Tuomilehto J, et al.: Comparison of the fasting and the 2-hour glucose criteria for diabetes in different Asian cohorts. Diabetologia 2000;43:1470–1475.
57. Resnick HE, Harris MI, Brock DB, Harris TB: American Diabetes Association diabetes diagnostic criteria, advancing age, and cardiovascular disease risk profiles. Results from the Third National Health and Nutrition Examination Survey. Diabetes Care 2000;23(2):176–180.
58. Pan X, Li G, Hu Y, et al.: Effects of diet and exercise in preventing NIDDM in people with impaired glucose tolerance: the Da Qing IGT and Diabetes Study. Diabetes Care 1997;20(4):537–544.
59. Tuomilehto J, Lindstrom J, Eriksson JG, et al.: Prevention of type 2 diabetes mellitus by changes in lifestyle among subjects with impaired glucose tolerance. N Engl J Med 2001; 344(18):1343–1350.
60. Knowler WC, Barrett-Connor E, Fowler SE, et al.: Reduction in the incidence of type 2 diabetes with lifestyle intervention or metformin. N Engl J Med 2002;346(6):393–403.
61. Chiasson JL, Josse RG, Gomis R, et al.: Acarbose for prevention of type 2 diabetes mellitus: the STOP-NIDDM randomised trial. Lancet 2002;359(9323):2072–2077.
62. Yang W, Lin L, Qi J, et al.: The preventive effect of acarbose and metformin on the progression to diabetes mellitus in the IGT population: a 3-year multicenter prospective study. Clin J Endocrinol Metab 2001;17(3):131–135.
63. Dowse GK, Zimmet PZ, King H: Relationship between prevalence of impaired glucose tolerance and NIDDM in a population. Diabetes Care 1991;14(11):968–974.
64. Coughlan A,McCarty DJ, Jorgensen LN, Zimmet P: The epidemicof NIDD Min Asian and Pacific Island populations: prevalence and risk factors. Horm Metab Res 1997;29(7):323–331.
65. Ramachandran A, Snehalatha C, Dharmaraj D, Viswanathan M: Prevalence of glucose intolerance in Asian Indians. Urban–rural difference and significance of upper body adiposity. Diabetes Care 1992;15(10):1348–1355.
66. Ramachandran A, Snehalatha C, Latha E, et al.: Rising prevalence of NIDDM in an urban population in India. Diabetologia 1997;40(2):232–237.

67. Ramachandran A, Snehalatha C,KapurA, et al.:High prevalence of diabetes and impaired glucose tolerance in India: National Urban Diabetes Survey. Diabetologia 2001;44:1094–1101.
68. National Diabetes Co-operative Study Group: A mass survey of diabetes mellitus in a population of 300 000 in 14 provinces and municipalities in China. Clin J Intern Med 1981;20(11):678–683.
69. Shanghai Diabetes Research Cooperative Group: A survey of diabetes mellitus among the population in Shanghai.Natl Med J Chin 1980;60:323–329.
70. Pan XR, Hu YH, Li GW, et al.: Impaired glucose tolerance and its relationship to ECG-indicated coronary heart disease and risk factors among Chinese: Da Qing IGT and Diabetes Study. Diabetes Care 1993;16(1):150–156.
71. Hu YH, Li GW, Pan XR: Incidence of NIDD Min Daqing and forecasting of NIDDM in China in 21st century. Zhonghua Nei Ke Za Zhi 1993;32(3):173–175.
72. Xiang HD, Wu W, Liu CQ, et al.: An epidemiological study on diabetes mellitus 1995–1996, in China. Chin J Diabetes 1998; 6(3):131–133.
73. Yang Z,Zheng H, Tong Z: Prevalence of diabetes and IGT in elderly population of Beijing in 1997. Chin J Geriatr 2001;20:290–293.
74. Jia WP, Xiang KS, Chen L, et al.: Epidemiological study on obesity and its comorbidities in urban Chinese older than 20 years of age in Shanghai, China. Obes Rev 2002;3(3):157–165.
75. Dong YH, Gao WG, Nan H, for the Qingdao Epidemiology Study Group: The prevalence of diabetes and the risk factor in Chinese population in Qingdao city [abstract]. In 18th International Diabetes Federation Congress, August 24–29, 2003, Paris, France, 2003.
76. Cockram CS,Woo J, Lau E, et al.: The prevalence of diabetes mellitus and impaired glucose tolerance among Hong Kong Chinese adults of working age. Diabetes Res Clin Pract 1993;21(1):67–73.
77. Janus ED, Watt NM, Lam KS, et al.: The prevalence of diabetes, association with cardiovascular risk factors and implications of diagnostic criteria (ADA 1997 and WHO 1998) in a 1996 community-based population study in Hong Kong Chinese. Diabet Med 2000;17(10):741–745.
78. Chen HD, Shaw CK, TsengWP, et al.: Prevalence of diabetes mellitus and impaired glucose tolerance in Aborigines and Chinese in eastern Taiwan. Diabetes Res Clin Pract 1997;38(3):199–205.
79. Midthjell K, Kruger O, Holmen J, et al.: Rapid changes in the prevalence of obesity and known diabetes in an adult Norwegian population. The Nord-Trondelag Health Surveys: 1984–1986 and 1995–1997. Diabetes Care 1999;22(11):1813–1820.
80. Drivsholm T, Ibsen H, Schroll M, et al.: Increasing prevalence of diabetes mellitus and impaired glucose tolerance among 60-year-old Danes. Diabet Med 2001;18:126–132.
81. Andersson DK, Svardsudd K, Tibblin G: Prevalence and incidence of diabetes in a Swedish community 1972–1987. Diabet Med 1991;8(5):428–434.
82. 82 Laakso M, Reunanen A, Klaukka T, et al.: Changes in the prevalence and incidence of diabetes mellitus in Finnish adults, 1970–1987. Am J Epidemiol 1991;133(9):850–857.
83. Gu K, Cowie CC, Harris MI: Diabetes and decline in heart disease mortality in US adults. JAMA 1999;281(14):1291–1297.
84. KnowlerWC,PettittDJ, SaadMF, Bennett PH: Diabetesmellitus in the Pima Indians: incidence, risk factors and pathogenesis. Diabetes Metab Rev 1990;6(1):1–27.
85. Burke JP,Williams K, Gaskill SP, et al.: Rapid rise in the incidence of type 2 diabetes from 1987 to 1996: results from the San Antonio Heart Study. Arch Intern Med 1999;159(13):1450–1456.
86. Savage PJ, Bennett PH, Senter RG, Miller M: High prevalence of diabetes in young Pima Indians: evidence of phenotypic variation in a genetically isolated population. Diabetes 1979; 28(10):937–942.
87. KnowlerWC, PettittDJ, Savage PJ, Bennett PH:Diabetes incidence in Pima Indians: contributions of obesity and parental diabetes. Am J Epidemiol 1981;113(2):144–156.
88. Dabelea D, Hanson RL, Bennett PH, et al.: Increasing prevalence of Type II diabetes in American Indian children. Diabetologia 1998;41(8):904–910.
89. Acton KJ, Burrows NR, Moore K, et al.: Trends in diabetes prevalence among American Indian and Alaska native children, adolescents, and young adults.AmJ PublicHealth 2002;92(9):1485–1490.
90. Burrows NR,Geiss LS,Engelgau MM,Acton KJ: Prevalence of diabetes among Native Americans and Alaska Natives, 1990–1997: an increasing burden. Diabetes Care 2000;23(12):1786–1790.

91. Harris SB, Perkins BA,Whalen-Brough E: Non-insulin-dependent diabetes mellitus among First Nations children. New entity among First Nations people of north-western Ontario. Can Fam Physician 1996;42:869–876.
92. Pinhas-Hamiel O, Dolan LM, Daniels SR, et al.: Increased incidence of non-insulin-dependent diabetes mellitus among adolescents. J Pediatr 1996;128 (5, pt 1): 608–615.
93. Kitagawa T, Owada M, Urakami T, Tajima N: Epidemiology of type 1 (insulin-dependent) and type 2 (non-insulin-dependent) diabetes mellitus in Japanese children. Diabetes Res Clin Pract 1994;24(Suppl):S7–S13.
94. Kitagawa T, Owada M, Urakami T, Yamauchi K: Increased incidence of non-insulin dependent diabetes mellitus among Japanese schoolchildren correlates with an increased intake of animal protein and fat. Clin Pediatr (Phila) 1998;37(2):111–115.
95. Owada M, Hanaoka Y, Tanimoto Y, Kitagawa T: Descriptive epidemiology of non-insulin dependent diabetes mellitus detected by urine glucose screening in school children in Japan. Acta Paediatr Jpn 1990;32(6):716–724.
96. Kadiki OA, Reddy MR, Marzouk AA: Incidence of insulindependent diabetes (IDDM) and non-insulin-dependent diabetes (NIDDM) (0–34 years at onset) in Benghazi, Libya. Diabetes Res Clin Pract 1996;32(3):165–173.
97. Likitmaskul S, Kiattisathavee P, Chaichanwatanakul K, et al.: Increasing prevalence of type 2 diabetes mellitus in Thai children and adolescents associated with increasing prevalence of obesity. J Pediatr Endocrinol Metab 2003;16(1):71–77.
98. Drake AJ, Smith A, Betts PR, et al.: Type 2 diabetes in obese white children. Arch Dis Child 2002;86(3):207–208.
99. Ehtisham S, Barrett TG, Shaw NJ: Type 2 diabetes mellitus in UK children—an emerging problem. Diabet Med 2000; 17(12):867–871.
100. Ehtisham S, Kirk J,McEvilly A, et al.: Prevalence of type 2 diabetes in children in Birmingham. BMJ 2001;322(7299):1428.
101. Ramachandran A, Snehalatha C, Satyavani K, et al.: Type 2 diabetes in Asian-Indian urban children. Diabetes Care 2003;26(4):1022–1025.
102. Wei JN, Chuang LM, Lin CC, et al.: Childhood diabetes identified in mass urine screening program in Taiwan, 1993–1999. Diabetes Res Clin Pract 2003;59(3):201–206.
103. Neufeld ND, Raffel LJ, Landon C, et al.: Early presentation of type 2 diabetes in Mexican-American youth. Diabetes Care 1998;21(1):80–86.
104. Willi SM, Kennedy A, Wojciechowski B, Garvey W: Insulin resistance and defective glucose-insulin coupling in ketosis-prone type 2 diabetes of African-American youth. Diabetes 1998;47(Suppl 1):A306.
105. Glaser NS, Jones KL: Non-insulin dependent diabetes mellitus in Mexican-American children. West J Med 1998;168(1):11–16.
106. Hale DE, Dabbey MM: Non-insulin dependent diabetes in Hispanic youth (type 2Y). Diabetes 1998;47(Suppl 1):A82.
107. Macaluso CJ, Bauer UE, Deeb LC, et al.: Type 2 diabetes mellitus among Florida children and adolescents, 1994 through 1998. Public Health Rep 2002;117:373–379.
108. Rosenbloom AL, Joe JR, Young RS, Winter WE: Emerging epidemic of type 2 diabetes in youth. Diabetes Care 1999;22(2): 345–354.
109. Fagot-Campagna A, Pettitt DJ, Engelgau MM, et al.: Type 2 diabetes among North American children and adolescents: an epidemiologic review and a public health perspective. J Pediatr 2000;136(5):664–672.
110. Freedman DS, Srinivasan SR, Burke GL, et al.: Relation of body fat distribution to hyperinsulinemia in children and adolescents: the Bogalusa Heart Study. Am J Clin Nutr 1987;46(3):403–410.
111. Arslanian S, Suprasongsin C: Insulin sensitivity, lipids, and body composition in childhood: is “syndrome X” present? J Clin Endocrinol Metab 1996;81(3):1058–1062.
112. Haffner SM, Stern MP, Mitchell BD, et al.: Incidence of type II diabetes in Mexican Americans predicted by fasting insulin and glucose levels, obesity, and body-fat distribution. Diabetes 1990;39(3):283–288.

113. Thomas-Dobersen DA, Butler-Simon N, Fleshner M: Evaluation of a weight management intervention program in adolescents with insulin-dependent diabetes mellitus. J Am Diet Assoc 1993; 93(5):535–540.
114. Pinhas-Hamiel O, Dolan LM, Zeitler PS: Diabetic ketoacidosis among obese African-American adolescents with NIDDM. Diabetes Care 1997;20(4):484–486.
115. Pihoker C, Scott CR, Lensing SY, et al.:Non-insulin dependent diabetes mellitus in African-American youths of Arkansas. Clin Pediatr (Phila) 1998;37(2):97–102.
116. Savola K, Laara E, Vahasalo P, et al.: Dynamic pattern of disease-associated autoantibodies in siblings of children with type 1 diabetes: a population-based study. Diabetes 2001;50(11): 2625–2632.
117. Kimpimaki T, Kulmala P, Savola K, et al.: Natural history of beta-cell autoimmunity in young children with increased genetic susceptibility to type 1 diabetes recruited from the general population. J Clin Endocrinol Metab 2002;87(10):4572–4579.
118. Ogden CL, Flegal KM, Carroll MD, Johnson CL: Prevalence and trends in overweight among US children and adolescents, 1999–2000. JAMA 2002;288(14):1728–1732.
119. Strauss RS, Pollack HA: Epidemic increase in childhood overweight, 1986–1998. JAMA 2001;286(22):2845–2848.
120. Amiel SA, Sherwin RS, Simonson DC, et al.: Impaired insulin action in puberty. A contributing factor to poor glycemic control in adolescents with diabetes. N Engl J Med 1986;315(4): 215–219.
121. Arslanian S, Suprasongsin C: Testosterone treatment in adolescents with delayed puberty: changes in body composition, protein, fat, and glucose metabolism. J Clin Endocrinol Metab 1997;82(10):3213–3220.
122. Caprio S, Plewe G, Diamond MP, et al.: Increased insulin secretion in puberty: a compensatory response to reductions in insulin sensitivity. J Pediatr 1989;114(6):963–977.
123. GoranMI, Gower BA: Longitudinal study on pubertal insulin resistance. Diabetes 2001;50(11):2444–2450.
124. Bloch CA, Clemons P, Sperling MA: Puberty decreases insulin sensitivity.J Pediatr 1987;110(3):481–487.
125. Dabelea D, Hanson RL, Lindsay RS, et al.: Intrauterine exposure to diabetes conveys risks for type 2 diabetes and obesity: a study of discordant sibships. Diabetes 2000;49(12):2208–2211.
126. Barker DJ, Martyn CN: The maternal and fetal origins of cardiovascular disease. J Epidemiol Community Health 1992;46(1):8–11.
127. Hales CN, Barker DJ, Clark PM, et al.: Fetal and infant growth and impaired glucose tolerance at age 64. BMJ 1991; 303(6809):1019–1022.
128. Young TK, Martens PJ, Taback SP, et al.: Type 2 diabetes mellitus in children: prenatal and early infancy risk factors among native Canadians. Arch Pediatr Adolesc Med 2002;156(7):651–655.
129. Pettitt D, Knowler WC: Long-term effects of the intrauterine environment, birth weight, and breast-feeding in Pima Indians. Diabetes Care 1998;21(Suppl 2):B138–B141.
130. Pettitt DJ, Forman MR, Hanson RL, et al.: Breastfeeding and incidence of non-insulin-dependent diabetes mellitus in Pima Indians. Lancet 1997; 350(9072):166–168.
131. Krakoff J, Lindsay RS, Looker HC, et al.: Incidence of retinopathy and nephropathy in youth-onset compared with adult-onset type 2 diabetes. Diabetes Care 2003;26(1):76–81.
132. Dean HJ, Flett B: Natural history of type 2 diabetes diagnosed in childhood: long term follow-up in young adult years. Diabetes 2002;51:A24–A25.
133. Ferrara A, Imperatore G, Quesenberry CP, et al.: Characteristics and quality of health care of type 1 (T1DM) and type 2 (T2DM) diabetic youths: the Northern California Kaiser Permanente (NCKP) Diabetes Registry. Diabetes 2002;51(Suppl 2):105.
134. Nelson RG, Sievers ML, Knowler WC, et al.: Low incidence of fatal coronary heart disease in Pima Indians despite high prevalence of non-insulin-dependent diabetes. Circulation 1990;81(3):987–995.
135. American Diabetes Association: Type 2 diabetes in children and adolescents. Diabetes Care 2000;23:381–389.
136. Freedman DS, Serdula MK, Percy CA, et al.: Obesity, levels of lipids and glucose, and smoking among Navajo adolescents. J Nutr 1997;127:2120S–2127S.
137. Dean HJ, Young TK, Flett B, Wood-Steiman P: Screening for type 2 diabetes in aboriginal children in northern Canada. Lancet 1998;352:1523–1524.

138. Fagot-Campagna A, Saadine JB, Fiegal KM, Beckles GL: Diabetes, impaired fasting glucose, and elevated HbA1c in US adolescents; the Third National Health and Nutrition Examination Survey. Diabetes Care 2001;24:834–837.
139. Scott CR, Smith JM, Creddock MM, Pihoker C: Characteristics of youth-onset noninsulin-dependent diabetes mellitus and insulin-dependent diabetes mellitus at diagnosis. Pediatrics 1997; 100:84–91.
140. O'Dea K: Obesity and diabetes in "the land of milk and honey." Diabetes Metab Rev 1992;8(4):373–388.
141. Knowler WC, Saad MF, Pettitt DJ, et al.: Determinants of diabetes mellitus in the Pima Indians. Diabetes Care 1993;16(1):216–227.
142. Haffner SM: Epidemiology of type 2 diabetes: risk factors. Diabetes Care 1998;21(Suppl 3):C3–C6.
143. Ford ES, Williamson DF, Liu S: Weight change and diabetes incidence: findings from a national cohort of US adults.AmJ Epidemiol 1997;146(3):214–222.
144. Colditz GA,Willett WC, Rotnitzky A,Manson JE:Weight gain as a risk factor for clinical diabetes mellitus in women. Ann Intern Med 1995;122(7):481–486.
145. Chang CJ, Wu CH, Lu FH, et al.: Discriminating glucose tolerance status by regions of interest of dual-energy X-ray absorptiometry. Clinical implications of body fat distribution. Diabetes Care 1999;22(12):1938–1943.
146. Goodpaster BH, Krishnaswami S, Resnick H, et al.: Association between regional adipose tissue distribution and both type 2 diabetes and impaired glucose tolerance in elderly men and women. Diabetes Care 2003;26(2):372–379.
147. Despres JP: Abdominal obesity as important component of insulin-resistance syndrome. Nutrition 1993;9(5):452–459.
148. Rush EC, Plank LD, Mitchelson E, Laulu MS: Central obesity and risk for type 2 diabetes in Maori, Pacific, and European young men in New Zealand. Food Nutr Bull 2002;23(3, suppl):82–86.
149. Hayashi T, Boyko EJ, Leonetti DL, et al.: Visceral adiposity and the risk of impaired glucose tolerance: a prospective study among Japanese Americans. Diabetes Care 2003;26(3):650–655.
150. Cruz ML, Bergman RN, Goran MI: Unique effect of visceral fat on insulin sensitivity in obese Hispanic children with a family history of type 2 diabetes. Diabetes Care 2002;25(9):1631–1636.
151. Hu FB, Sigal RJ, Rich-Edwards JW, et al.:Walking compared with vigorous physical activity and risk of type 2 diabetes in women: a prospective study. JAMA 1999;282(15):1433–1439.
152. Mayer-Davis EJ, D'Agostino R Jr, Karter AJ, et al.: Intensity and amount of physical activity in relation to insulin sensitivity: the Insulin Resistance Atherosclerosis Study. JAMA 1998;279 (9):669–674.
153. Kriska AM, Pereira MA, Hanson RL, et al.: Association of physical activity and serum insulin concentrations in two populations at high risk for type 2 diabetes but differing by BMI 2001; 24(7):1175–1180.
154. Manson JE, Nathan DM, Krolewski AS, et al.: A prospective study of exercise and incidence of diabetes among US male physicians. JAMA 1992;268(1):63–67.
155. Helmrich SP, Ragland DR, Leung RW, Paffenbarger RS Jr. Physical activity and reduced occurrence of non-insulin-dependent diabetes mellitus. N Engl J Med 1991;325(3):147–152.
156. Manson JE, Rimm EB, Stampfer MJ, et al.: Physical activity and incidence of non-insulin-dependent diabetes mellitus in women. Lancet 1991;338(8770):774–778.
157. Schranz A, Tuomilehto J, Marti B, et al.: Low physical activity and worsening of glucose tolerance: results from a 2-year follow-up of a population sample in Malta. Diabetes Res Clin Pract 1991;11:127–136.
158. Burchfiel CM, Sharp DS, Curb JD, et al.: Physical activity and incidence of diabetes: the Honolulu Heart Program. Am J Epidemiol 1995;141(4):360–368.
159. Lynch J,Helmrich SP, Lakka TA, et al.:Moderately intense physical activities and high levels of cardiorespiratory fitness reduce the risk of non-insulin-dependent diabetes mellitus in middle-aged men. Arch Intern Med 1996;156(12):1307–1314.
160. Haapanen N, Miilunpalo S, Vuori I, et al.: Association of leisure time physical activity with the risk of coronary heart disease, hypertension and diabetes in middle-aged men and women. Int J Epidemiol 1997;26(4):739–747.

161. Folsom AR, Kushi LH, Hong CP: Physical activity and incident diabetes mellitus in postmenopausal women. Am J Public Health 2000;90(1):134–138.
162. Hu FB, Leitzmann MF, Stampfer MJ, et al.: Physical activity and television watching in relation to risk for type 2 diabetes mellitus in men. Arch Intern Med 2001;161(12):1542–1548.
163. Hu FB, Li TY, Colditz GA, et al.: Television watching and other sedentary behaviors in relation to risk of obesity and type 2 diabetes mellitus in women. JAMA 2003;289(14):1785–1791.
164. Eriksson KF, Lindgarde F: Prevention of type 2 (non-insulindependent) diabetes mellitus by diet and physical exercise. The 6-yearMalmö feasibility study. Diabetologia 1991;34(12):891–898.
165. Danforth E Jr. Diet and obesity. Am J Clin Nutr 1985;41(5, suppl):1132–1145.
166. Storlien LH, James DE, Burleigh KM, et al.: Fat feeding causes widespread in vivo insulin resistance, decreased energy expenditure, and obesity in rats. Am J Physiol 1986;251(5, pt 1):E576–E583.
167. Storlien LH, Pan DA, Kriketos AD, Baur LA:High fat diet-induced insulin resistance. Lessons and implications from animal studies. Ann N Y Acad Sci 1993;683:82–90.
168. Kawate R, Yamakido M, Nishimoto Y, et al.: Diabetes mellitus and its vascular complications in Japanese migrants on the Island of Hawaii. Diabetes Care 1979;2(2):161–170.
169. Stern MP, Gonzalez C, Mitchell BD, et al.: Genetic and environmental determinants of type II diabetes in Mexico City and San Antonio. Diabetes 1992;41(4):484–492.
170. Tsunehara CH, Leonetti DL, Fujimoto WY: Diet of second generation Japanese-American men with and without non-insulin dependent diabetes. Am J Clin Nutr 1990;52(4):731–738.
171. Marshall JA, Hamman RF, Baxter J: High-fat, low-carbohydrate diet and the etiology of non-insulin-dependent diabetes mellitus: the San Luis Valley Diabetes Study. Am J Epidemiol 1991;134 (6):590–603.
172. Feskens EJ, Virtanen SM, Rasanen L, et al.: Dietary factors determining diabetes and impaired glucose tolerance. A 20-year follow-up of the Finnish and Dutch cohorts of the seven countries study. Diabetes Care 1995;18(8):1104–1112.
173. Colditz GA,Manson JE, Stampfer MJ, et al.: Diet and risk of clinical diabetes in women. Am J Clin Nutr 1992;55(5):1018–1023.
174. Lundgren H, Bengtsson C, Blohme G, et al.: Dietary habits and incidence of noninsulin-dependent diabetes mellitus in a population study of women in Gotnenburg, Sweden. Am J Clin Nutr 198 9;49:708–712.
175. Virtanen SM, Rasanen L, Maenpaa J, Akerblom HK: Dietary survey of Finnish adolescent diabetics and non-diabetic controls. Acta Paediatr Scand 1987;76(5):801–808.
176. Marshall JA, Bessesen DH, Hamman RF: High saturated fat and low starch and fibre are associated with hyper-insulinaemia in a non-diabetic population: the San Luis Valley Diabetes Study. Diabetologia 1997;40(4):430–438.
177. Adler AI, Boyko EJ, Schraer CD, Murphy NJ: Lower prevalence of impaired glucose tolerance and diabetes associated with daily seal oil or salmon consumption among Alaska Natives. Diabetes Care 1994;17(12):1498–1501.
178. Vessby B: Effects of omega 3 fatty acids on glucose and lipid metabolism in non-insulin-dependent diabetes mellitus. World Rev Nutr Diet 1991;66:407–416.
179. Storlien LH, Kraegen EW, Chisholm DJ, et al.: Fish oil prevents insulin resistance induced by high-fat feeding in rats. Science 1987;237(4817):885–888.
180. Hu FB, van Dam RM, Liu S: Diet and risk of Type II diabetes: the role of types of fat and carbohydrate. Diabetologia 2001;44(7):805–817.
181. Feskens EJ, Bowles CH, Kromhout D: Carbohydrate in-take and body mass index in relation to the risk of glucose intolerance in an elderly population. Am J Clin Nutr 1991;54(1):136–140.
182. Wolever TM, Hamad S, Gittelsohn J, et al.: Low dietary fiber and high protein intakes associated with newly diagnosed diabetes in a remote aboriginal community. Am J Clin Nutr 1997; 66(6):1470–1474.
183. Jenkins DJ, Wolever TM,Taylor RH, et al.: Glycemic index of foods: a physiological basis for carbohydrate exchange. Am J Clin Nutr 1981;34(3):362–366.
184. Salmeron J, Ascherio A, Rimm EB, et al.: Dietary fiber, glycemic load, and risk of NIDDM in men. Diabetes Care 1997;20(4):545–550.
185. Meyer KA, Kushi LH, Jacobs DR Jr, et al.: Carbohydrates, dietary fiber, and incident type 2 diabetes in older women. Am J Clin Nutr 2000;71(4):921–930.

186. van Dam RM, Visscher AW, Feskens EJ, et al.: Dietary glycemic index in relation to metabolic risk factors and incidence of coronary heart disease: the Zutphen Elderly Study. Eur J Clin Nutr 2000;54(9):726–731.
187. Trowell HC: Dietary-fiber hypothesis of the etiology of diabetes mellitus. Diabetes 1975;24(8):762–765.
188. Burkitt DP, Trowell HC: Dietary fibre and Western diseases. Ir Med J 1977;70(9):272–277.
189. Salmeron J,Manson JE, Stampfer MJ, et al.: Dietary fiber, glycemic load, and risk of non-insulin-dependent diabetes mellitus in women. JAMA 1997;277(6):472–477.
190. Fung TT, Hu FB, Pereira MA, et al.: Whole-grain intake and the risk of type 2 diabetes: a prospective study in men. Am J Clin Nutr 2002;76(3):535–540.
191. Bakker SJ, Hoogeveen EK, Nijpels G, et al.: The association of dietary fibres with glucose tolerance is partly explained by concomitant intake of thiamine: the Hoorn Study. Diabetologia 1998;41(10):1168–1175.
192. Jenkins DJ, Axelsen M, Kendall CW, et al.: Dietary fibre, lente carbohydrates and the insulin-resistant diseases. Br J Nutr 2000;83(Suppl 1):S157–S163.
193. Williams DE, Wareham NJ, Cox BD, et al.: Frequent salad vegetable consumption is associated with a reduction in the risk of diabetes mellitus. J Clin Epidemiol 1999;52(4):329–335.
194. Ylonen K, Alfthan G, Groop L, et al.: Dietary intakes and plasma concentrations of carotenoids and tocopherols in relation to glucose metabolism in subjects at high risk of type 2 diabetes: the Botnia Dietary Study. Am J Clin Nutr 2003;77(6):1434–1441.
195. Ford ES, Mokdad AH, Giles WH, Brown DW: The metabolic syndrome and antioxidant concentrations: findings from the Third National Health and Nutrition Examination Survey. Diabetes 2003;52(9):2346–2352.
196. Sargeant LA, Khaw KT, Bingham S, et al.: Cigarette smoking and glycaemia: the EPIC-Norfolk Study. Int J Epidemiol 2001;30(3):547–554.
197. Williams DE, Prevost AT, Whichelow MJ, et al.: A cross-sectional study of dietary patterns with glucose intolerance and other features of the metabolic syndrome. Br J Nutr 2000;83(3):257–266.
198. van Dam RM, Rimm EB, Willett WC, et al.: Dietary patterns and risk for type 2 diabetes mellitus in U.S. men. Ann Intern Med 2002;136(3):201–209.
199. Williams DE, Knowler WC, Smith CJ, et al.: The effect of Indian or Anglo dietary preference on the incidence of diabetes in Pima Indians. Diabetes Care 2001;24(5):811–816.
200. Brand JC, Snow BJ, Nabhan GP, Truswell AS: Plasma glucose and insulin responses to traditional Pima Indian meals. Am J Clin Nutr 1990;51(3):416–420.
201. Thorburn AW, Brand JC, O'Dea K, et al.: Plasma glucose and insulin responses to starchy foods in Australian aborigines: a population now at high risk of diabetes. Am J Clin Nutr 1987;46(2):282–285.
202. Jarrett RJ, Shipley MJ: Type 2 (non-insulin-dependent) diabetes mellitus and cardiovascular disease—putative association via common antecedents; further evidence from the Whitehall Study. Diabetologia 1988;31(10):737–740.
203. Stern MP: Diabetes and cardiovascular disease.The "common soil" hypothesis. Diabetes 1995;44(4):369–374.
204. Yudkin JS, Stehouwer CD, Emeis JJ, Coppack SW: C-reactive protein in healthy subjects: associations with obesity, insulin resistance, and endothelial dysfunction: a potential role for cytokines originating from adipose tissue? Arterioscler Thromb Vasc Biol 1999;19(4):972–978.
205. Festa A, D'Agostino R Jr, Howard G, et al.: Chronic subclinical inflammation as part of the insulin resistance syndrome: the Insulin Resistance Atherosclerosis Study (IRAS). Circulation 2000;102(1):42–47.
206. Forouhi NG, Sattar N, McKeigue PM: Relation of C-reactive protein to body fat distribution and features of the metabolic syndrome in Europeans and South Asians. Int J Obes Relat Metab Disord 2001;25(9):1327–1331.
207. Pradhan AD, Manson JE, Rifai N, et al.: C-reactive protein, interleukin 6, and risk of developing type 2 diabetes mellitus. JAMA 2001;286(3):327–334.
208. Barzilay JI, Abraham L, Heckbert SR, et al.: The relation of markers of inflammation to the development of glucose disorders in the elderly: the Cardiovascular Health Study. Diabetes 2001;50(10):2384–2389.

209. Freeman DJ, Norrie J, Caslake MJ, et al.: C-reactive protein is an independent predictor of risk for the development of diabetes in the West of Scotland Coronary Prevention Study. Diabetes 2002;51(5):1596–600.
210. Festa A, D'Agostino R Jr, Tracy RP, Haffner SM: Elevated levels of acute-phase proteins and plasminogen activator inhibitor-1 predict the development of type 2 diabetes: the insulin resistance atherosclerosis study. Diabetes 2002;51(4):1131–1137.
211. Han TS, Sattar N,Williams K, et al.: Prospective study of C-reactive protein in relation to the development of diabetes and metabolic syndrome in the Mexico City Diabetes Study. Diabetes Care 2002;25(11):2016–2221.
212. Thorand B, Lowel H, Schneider A, et al.: C-reactive protein as a predictor for incident diabetes mellitus among middle-aged men: results from the MONICA Augsburg Cohort Study, 1984–1998. Arch Intern Med 2003;163(1):93–99.
213. Hundal RS, Petersen KF,Mayerson AB, et al.:Mechanism by which high-dose aspirin improves glucose metabolism in type 2 diabetes. J Clin Invest 2002;109(10):1321–1326.
214. Rimm EB, Manson JE, Stampfer MJ, et al.: Cigarette smoking and the risk of diabetes in women. Am J Public Health 1993;83(2):211–214.
215. Rimm EB, Chan J, Stampfer MJ, et al.: Prospective study of cigarette smoking, alcohol use, and the risk of diabetes in men. BMJ 1995;310(6979):555–559.
216. Will JC, Galuska DA, Ford ES, et al.: Cigarette smoking and diabetes mellitus: evidence of a positive association from a large prospective cohort study. Int J Epidemiol 2001;30(3):540–546.
217. Facchini FS, Hollenbeck CB, Jeppesen J, et al.: Insulin resistance and cigarette smoking. Lancet 1992;339(8802):1128–1130.
218. Attvall S, Fowelin J, Lager I, et al.: Smoking induces insulin resistance—a potential link with the insulin resistance syndrome. J InternMed 1993;233(4):327–332.