

## International Textbook of Diabetes Mellitus, 4th Ed., Chapter 6 - Insulin Gene Expression and Biosynthesis

### References:

1. Le Roith D, Hendricks SA, Lesniak MA, et al.: Insulin in brain and other extrapancreatic tissues of vertebrates and nonvertebrates. *Advances in Metabolic Disorders* 1983;10:303–340.
2. Smith KM, Olson DC, Hirose R, Hanahan D: Pancreatic gene expression in rare cells of thymic medulla: evidence for functional contribution to T cell tolerance. *International Immunology* 1997; 9(9):1355 – 1365.
3. Giddings SJ, Carnaghi L: Rat insulin II gene expression by extraplacental membranes. *A non-pancreatic source for fetal insulin. The Journal of Biological Chemistry* 1989;264(16):9462–9469.
4. Soares MB, Schon E, Henderson A, et al.: RNA-mediated gene duplication: the rat preproinsulin I gene is a functional retroposon. *Molecular and Cellular Biology* 1985;5(8):2090–2103.
5. Oliver-Krasinski JM, Stoffers DA: On the origin of the beta cell. *Genes & Development* 2008;22(15):1998 – 2021.
6. Andrali SS, Sampley ML, Vanderford NL, Ozcan S: Glucose regulation of insulin gene expression in pancreatic beta-cells. *The Biochemical Journal* 2008;415(1):1–10.
7. Ban N, Yamada Y, Someya Y, et al.: Activating transcription factor-2 is a positive regulator in CaM kinase IV-induced human insulin gene expression. *Diabetes* 2000;49(7):1142–1148.
8. Lawrence MC, Bhatt HS, Watterson JM, Easom RA: Regulation of insulin gene transcription by a Ca(2+)-responsive pathway involving calcineurin and nuclear factor of activated T cells. *Molecular Endocrinology* 2001;15(10):1758–1767.
9. Lawrence MC, Shao C, McGlynn K, et al.: Multiple chromatin-bound protein kinases assemble factors that regulate insulin gene transcription. *PNAS* 2009;106(52):22181–22186.
10. Lu M, Seufert J, Habener JF: Pancreatic beta-cell-specific repression of insulin gene transcription by CCAAT/enhancer-binding protein beta. Inhibitory interactions with basic helix-loop-helix transcription factor E47. *The Journal of Biological Chemistry* 1997; 272(45):28349 – 28359.
11. Inagaki N, Maekawa T, Sudo T, et al.: c-Jun represses the human insulin promoter activity that depends on multiple cAMP response elements. *PNAS* 1992;89(3):1045 – 1049.
12. Heit JJ: Calcineurin/NFAT signaling in the beta-cell: from diabetes to new therapeutics. *BioEssays : news and reviews in molecular, cellular and developmental biology* 2007;29(10):1011 – 1021.
13. Qiu Y, Guo M, Huang S, Stein R: Insulin gene transcription is mediated by interactions between the p300 coactivator and PDX-1, BETA2, and E47. *Molecular and Cellular Biology* 2002;22(2): 412 – 420.
14. Stein R: Insulin gene transcription: the factors involved in cell-type specific and glucose-regulated expression in islet beta-cells are also essential

- during pancreatic development. In: Cherrington A, Jefferson J (eds.) *Handbook of Physiology, Section 7: The Endocrine System*. Washington, DC: American Physiology Society, 2001, pp. 25 – 78.
15. Ohneda K, Mirmira RG, Wang J, et al.: The homeodomain of PDX-1 mediates multiple protein-protein interactions in the formation of a transcriptional activation complex on the insulin promoter. *Molecular and Cellular Biology* 2000;20(3):900 – 911.
  16. Stoffers DA, Thomas MK, Habener JF: Homeodomain protein IDX-1. *A master regulator of pancreas development and insulin gene expression. Trends in Endocrinology & Metabolism* 1997;8:145 – 151.
  17. Jonsson J, Ahlgren U, Edlund T, Edlund H: IPF-1, a homeodomain protein with dual function in pancreas development. *International Journal of Developmental Biology* 1995;39:789 – 798.
  18. Wang H, Maechler P, Ritz-Laser B, et al.: Pdx1 level defines pancreatic gene expression pattern and cell lineage differentiation. *The Journal of Biological Chemistry* 2001;276(27):25279 – 25286.
  19. Naya FJ, Huang H-P, Qiu Y, et al.: Diabetes, defective pancreatic morphogenesis, and abnormal enteroendocrine differentiation in BETA2/NeuroD-deficient mice. *Genes and Development* 1997;11: 2323 – 2334.
  20. Gu C, Stein GH, Pan N, et al.: Pancreatic beta cells require NeuroD to achieve and maintain functional maturity. *Cell Metabolism* 2010; 11(4):298 – 310.
  21. Malecki MT, Jhala US, Antonellis A, et al.: Mutations in NEUROD1 are associated with the development of type 2 diabetes mellitus. *Nature Genetics* 1999;23(3):323 – 328.
  22. Artner I, Le Lay J, Hang Y, et al.: MafB: an activator of the glucagon gene expressed in developing islet alpha- and beta-cells. *Diabetes* 2006;55(2):297 – 304.
  23. Nishimura W, Kondo T, Salameh T, et al.: A switch from MafB to MafA expression accompanies differentiation to pancreatic beta-cells. *Developmental Biology* 2006;293(2):526 – 539.
  24. Matsuoka TA, Artner I, Henderson E, et al.: The MafA transcription factor appears to be responsible for tissue-specific expression of insulin. *PNAS* 2004;101(9):2930 – 2933.
  25. Dai C, Brissova M, Hang Y, et al.: Islet-enriched gene expression and glucose-induced insulin secretion in human islets. *Diabetologia* 2012;55(3):707 – 718.
  26. Zhang C, Moriguchi T, Kajihara M, et al.: MafA is a key regulator of glucose-stimulated insulin secretion. *Molecular and Cellular Biology* 2005;25(12):4969 – 4976.
  27. Artner I, Hang Y, Mazur M, et al.: MafA and MafB regulate genes critical to {beta}-cells in a unique temporal manner. *Diabetes* 2010;59(10):2530 – 2539.
  28. Mosley AL, Ozcan S: The pancreatic duodenal homeobox-1 protein (Pdx-1) interacts with histone deacetylases Hdac-1 and Hdac-2 on low levels of

- glucose. *The Journal of Biological Chemistry* 2004;279(52):54241 – 54247.
29. Mosley AL, Ozcan S: Glucose regulates insulin gene transcription by hyperacetylation of histone h4. *The Journal of Biological Chemistry* 2003;278(22):19660 – 19666.
  30. Kishi A, Nakamura T, Nishio Y, et al.: Sumoylation of Pdx1 is associated with its nuclear localization and insulin gene activation. *American Journal of Physiology - Endocrinology and Metabolism* 2003; 284(4):E830 – 840.
  31. Gao Y, Miyazaki J, Hart GW: The transcription factor PDX-1 is post-translationally modified by O-linked N-acetylglucosamine and this modification is correlated with its DNA binding activity and insulin secretion in min6 beta-cells. *Archives of Biochemistry and Biophysics* 2003;415(2):155 – 163.
  32. Kebede M, Ferdaoussi M, Mancini A, et al.: Glucose activates free fatty acid receptor 1 gene transcription via phosphatidylinositol-3- kinase-dependent O-GlcNAcylation of pancreas-duodenum homeobox-1. *PNAS* 2012;109(7):2376 – 2381.
  33. Khoo S, Griffen SC, Xia Y, et al.: Regulation of insulin gene transcription by ERK1 and ERK2 in pancreatic beta cells. *The Journal of Biological Chemistry* 2003;278(35):32969 – 32977.
  34. Andrali SS, Qian Q, Ozcan S: Glucose mediates the translocation of NeuroD1 by O-linked glycosylation. *The Journal of Biological Chemistry* 2007;282(21):15589 – 15596.
  35. Guo S, Vanderford NL, Stein R: Phosphorylation within the MafA N-terminus regulates C-terminal dimerization and DNA binding. *Journal of Biological Chemistry* 2010;285(17):12655 – 12661.
  36. Rocques N, Abou Zeid N, Sii-Felice K, et al.: GSK-3-mediated phosphorylation enhances Maf-transforming activity. *Molecular Cell* 2007;28(4):584 – 597.
  37. Fred RG, Welsh N: The importance of RNA binding proteins in pre-proinsulin mRNA stability. *Molecular and Cellular Endocrinology [Review]* 2009;297(1 – 2):28 – 33.
  38. Giddings SJ, Chirgwin J, Permutt MA: The effects of fasting and feeding on preproinsulin messenger RNA in rats. *Journal of Clinical Investigation* 1981;67(4):952 – 960.
  39. Giddings SJ, Chirgwin J, Permutt MA: Effects of glucose on proinsulin messenger RNA in rats in vivo. *Diabetes* 1982;31(7):624 – 629.
  40. Shalwitz RA, Herbst T, Carnaghi LR, Giddings SJ: Time course for effects of hypoglycemia on insulin gene transcription in vivo. *Diabetes* 1994;43:929 – 934.
  41. Pipeleers DG: Heterogeneity in pancreatic  $\beta$ -cell population. *Diabetes* 1992;41:777 – 781.
  42. Moitoso de Vargas L, Sobolewski J, Siegel R, Moss LG: Individual  $\beta$ -cells within the intact islet differentially respond to glucose. *The Journal of Biological Chemistry* 1997;272:26573 – 26577.

43. Drucker DJ: The biology of incretin hormones. *Cell Metabolism* 2006;3(3):153 – 165.
44. Lamont BJ, Li Y, Kwan E, et al.: Pancreatic GLP-1 receptor activation is sufficient for incretin control of glucose metabolism in mice. *Journal of Clinical Investigation* 2012;122(1):388 – 402.
45. Smith PA, Sakura H, Coles B, et al.: Electrogenic arginine transport mediates stimulus-secretion coupling in mouse pancreatic  $\beta$ -cells. *Journal of Physiology* 1997;499:625 – 635.
46. Hussain MA, Habener JF: Glucagon-like peptide 1 increases glucose-dependent activity of the homeoprotein IDX-1 transactivating domain in pancreatic beta-cells. *Biochemical and Biophysical Research Communications* 2000;274(3):616 – 619.
47. Wang X, Cahill CM, Pineyro MA, et al.: Glucagon-like peptide-1 regulates the beta cell transcription factor, PDX-1, in insulinoma cells. *Endocrinology* 1999;140(10):4904 – 4907.
48. Lawrence MC, Bhatt HS, Easom RA: NFAT regulates insulin gene promoter activity in response to synergistic pathways induced by glucose and glucagon-like peptide-1. *Diabetes* 2002;51(3):691 – 698.
49. Winzell MS, Ahren B: G-protein-coupled receptors and islet function-implications for treatment of type 2 diabetes. *Pharmacology & Therapeutics* 2007;116(3):437–448.
50. Zhang HJ, Redmond JB, Andresen JM, Robertson RP: Somatostatin and epinephrine decrease insulin messenger ribonucleic acid in HIT cells through a pertussis toxin-sensitive mechanism. *Endocrinology* 1991;129(5):2409 – 2414.
51. Philippe J: Somatostatin inhibits insulin-gene expression through a posttranscriptional mechanism in a hamster islet cell line. *Diabetes* 1993;42:244 – 249.
52. Hussain MA, Daniel PB, Habener JF: Glucagon stimulates expression of the inducible cAMP early repressor and suppresses insulin gene expression in pancreatic beta-cells. *Diabetes* 2000;49(10):1681 – 1690.
53. Seufert J, Kieffer TJ, Habener JF: Leptin inhibits insulin gene transcription and reverses hyperinsulinemia in leptin-deficient ob/ob mice. *PNAS* 1999;96(2):674–679.
54. Leibiger IB, Leibiger B, Berggren P-O: Insulin signaling in the pancreatic  $\beta$ -cell. *Annual Review of Nutrition* [Review] 2008;28: 233 – 251.
55. Rhodes CJ, White MF, Leahy JL, Kahn SE: Direct autocrine action of insulin on  $\beta$ -cells: Does it make physiological sense? *Diabetes* 2013;62(7):2157 – 2163.
56. Anjos S, Polychronakos C: Mechanisms of genetic susceptibility to type I diabetes: beyond HLA. *Molecular Genetics and Metabolism* [Review] 2004;81(3):187–195.
57. Robertson RP, Harmon J, Tran PO, et al.: Glucose toxicity in beta-cells: type 2 diabetes, good radicals gone bad, and the glutathione connection. *Diabetes* 2003;52(3):581–587.

58. Poitout V, Robertson RP: Glucolipotoxicity: fuel excess and beta-cell dysfunction. *Endocrine Reviews* 2008;29(3):351–366.
59. Lu M, Seufert J, Habener JF: Pancreatic  $\beta$ -cell-specific repression of insulin gene transcription by CCAAT/Enhancer-binding protein  $\beta$ . Inhibitory interactions with basic helix-loop-helix transcription factor E47. *Journal of Biological Chemistry* 1997;272:28349–28359.
60. Lawrence MC, McGlynn K, Park BH, Cobb MH: ERK1/2-dependent activation of transcription factors required for acute and chronic effects of glucose on the insulin gene promoter. *The Journal of Biological Chemistry* 2005;280(29):26751–26759.
61. Robertson RP, Harmon JS: Diabetes, glucose toxicity, and oxidative stress: a case of double jeopardy for the pancreatic islet beta cell. *Free Radical Biology & Medicine* 2006;41(2):177–184.
62. Puddu A, Storace D, Odetti P, Viviani GL: Advanced glycation end-products affect transcription factors regulating insulin gene expression. *Biochemical and Biophysical Research Communications* 2010;395(1):122 – 125.
63. Tanaka Y, Gleason CE, Tran POT, et al.: Prevention of glucose toxicity in HIT-T15 cells and Zucker diabetic fatty rats by antioxidants. *PNAS* 1999;96:10857 – 10862.
64. Harmon JS, Bogdani M, Parazzoli SD, et al.: beta-Cell-specific over-expression of glutathione peroxidase preserves intranuclear MafA and reverses diabetes in db/db mice. *Endocrinology* 2009;150(11): 4855 – 4862.
65. Kaneto H, Xu G, Fujii N, et al.: Involvement of c-Jun N-terminal kinase in oxidative stress-mediated suppression of insulin gene expression. *The Journal of Biological Chemistry* 2002;277(33): 30010 – 30018.
66. Weir GC, Sharma A, Zangen DH, Bonner-Weir S: Transcription factor abnormalities as a cause of beta cell dysfunction in diabetes: a hypothesis. *Acta Diabetologica* 1997;34:177–184.
67. Laybutt DR, Sharma A, Sgroi DC, et al.: Genetic regulation of metabolic pathways in beta-cells disrupted by hyperglycemia. *The Journal of Biological Chemistry* 2002;277(13):10912–10921.
68. Jonas JC, Sharma A, Hasenkamp W, et al.: Chronic hyperglycemia triggers loss of pancreatic beta cell differentiation in an animal model of diabetes. *The Journal of Biological Chemistry* 1999;274(20):14112 – 14121.
69. Jonas JC, Laybutt DR, Steil GM, et al.: High glucose stimulates early response gene c-Myc expression in rat pancreatic beta cells. *The Journal of Biological Chemistry* 2001;276(38):35375–35381.
70. Laybutt DR, Weir GC, Kaneto H, et al.: Overexpression of c-Myc in beta-cells of transgenic mice causes proliferation and apoptosis, downregulation of insulin gene expression, and diabetes. *Diabetes* 2002;51(6):1793 – 1804.
71. Kaneto H, Sharma A, Suzuma K, et al.: Induction of c-Myc expression suppresses insulin gene transcription by inhibiting NeuroD/BETA2-

- mediated transcriptional activation. *The Journal of Biological Chemistry* 2002;277(15):12998–13006.
72. Mooradian AD, Haas MJ: Glucose-induced endoplasmic reticulum stress is independent of oxidative stress: a mechanistic explanation for the failure of antioxidant therapy in diabetes. *Free Radical Biology & Medicine* 2011;50(9):1140–1143.
  73. Tang C, Koulajian K, Schuikil, et al.: Glucose-induced beta cell dysfunction in vivo in rats: link between oxidative stress and endoplasmic reticulum stress. *Diabetologia* 2012;55(5):1366–1379.
  74. Unger RH: Lipotoxicity in the pathogenesis of obesity-dependent NIDDM. Genetic and clinical implications. *Diabetes* 1995;44: 863 – 870.
  75. Prentki M, Joly E, El-Assaad W, Roduit R: Malonyl-CoA signaling, lipid partitioning, and glucolipotoxicity: role in beta-cell adaptation and failure in the etiology of diabetes. *Diabetes* 2002;51(Suppl 3): S405 – 413.
  76. Kelle CL, Moore PC, Parazzoli SD, et al.: Palmitate inhibition of insulin gene expression is mediated at the transcriptional level via ceramide synthesis. *The Journal of Biological Chemistry* 2003; 278(32):30015 – 30021.
  77. Hagman DK, Parazzoli SD, Poitout V: Prolonged exposure of isolated rat islets to palmitate reduces PDX-1 and MafA binding to the insulin gene promoter. *Diabetes* 2005;54:A399-A.
  78. Hagman DK, Latour MG, Chakrabarti SK, et al.: Cyclical and alternating infusions of glucose and intralipid in rats inhibit insulin gene expression and Pdx-1 binding in islets. *Diabetes* 2008;57(2):424 – 431.
  79. Fontes G, Zarrouki B, Hagman DK, et al.: Glucolipotoxicity age-dependently impairs beta cell function in rats despite a marked increase in beta cell mass. *Diabetologia* 2010;53(11):2369 – 2379.
  80. Harmon JS, Gleason CE, Tanaka Y, et al.: In vivo prevention of hyperglycemia also prevents glucotoxic effects on PDX-1 and insulin gene expression. *Diabetes* 1999;48(10):1995 – 2000.
  81. da Silva Xavier G, Rutter J, Rutter GA: Involvement of Per-Arnt-Sim (PAS) kinase in the stimulation of preproinsulin and pancreatic duodenum homeobox 1 gene expression by glucose. *PNAS* 2004; 101(22):8319 – 8324.
  82. Fontes G, Semache M, Hagman DK, et al.: Involvement of Per-Arnt- Sim Kinase and extracellular-regulated kinases-1/2 in palmitate inhibition of insulin gene expression in pancreatic beta-cells. *Diabetes* 2009;58(9):2048 – 2058.
  83. Humphrey RK, Yu SM, Flores LE, Jhala US: Glucose regulates steady-state levels of PDX1 via the reciprocal actions of GSK3 and AKT kinases. *The Journal of Biological Chemistry* 2010; 285(5):3406 – 3416.
  84. Solinas G, Naugler W, Galimi F, et al.: Saturated fatty acids inhibit induction of insulin gene transcription by JNK-mediated phosphorylation of insulin-receptor substrates. *PNAS* 2006;103(44): 16454 – 16459.

85. Del Guerra S, Lupi R, Marselli L, et al.: Functional and molecular defects of pancreatic islets in human type 2 diabetes. *Diabetes* 2005; 54(3):727 – 735.
86. Butler AE, Robertson RP, Hernandez R, et al.: Beta cell nuclear musculoaponeurotic fibrosarcoma oncogene family A (MafA) is deficient in type 2 diabetes. *Diabetologia* 2012;55(11):2985 – 2988.
87. Yang BT, Dayeh TA, Kirkpatrick CL, et al.: Insulin promoter DNA methylation correlates negatively with insulin gene expression and positively with HbA(1c) levels in human pancreatic islets. *Diabetologia* 2011;54(2):360 – 367.
88. Steiner DF, Rhodes CJ, Philipson LH, et al.: Biosynthesis, processing and secretion of islet hormones: insulin, islet amyloid polypeptide (amylin), glucagon, somatostatin and pancreatic polypeptide. In: Jameson JL, De Groot LJ, de Kretser D, et al. (eds.) *Endocrinology*, 6th Edition, Adult and Pediatric, Vol. 1. Philadelphia, PA: Saunders/Elsevier, 2010, pp. 603 – 623.
89. Kong J, Lasko P: Translational control in cellular and developmental processes. *Nature Reviews Genetics* [Review] 2012;13(6):383 – 394.
90. Shan SO, Walter P: Co-translational protein targeting by the signal recognition particle. *FEBS Letters* [Review] 2005;579(4):921 – 926.
91. Rhodes CJ: Processing the insulin molecule. In: LeRoith D, Taylor SI, Olefsky JM (eds.). *Diabetes Mellitus: A Fundamental and Clinical Text*, 3rd edn. Philadelphia, PA: Lippincott Williams & Wilkins, 2004, pp. 27 – 50.
92. Uchizono Y, Alarcon C, Wicksteed BL, et al.: The balance between proinsulin biosynthesis and insulin secretion: where can imbalance lead? *Diabetes, Obesity and Metabolism* 2007;9 (Suppl 2):56 – 66.
93. Orci L: The insulin factory: a tour of the plant surroundings and a visit to the assembly line. *Diabetologia* 1985;28:528 – 546.
94. Marsh BJ, Volkmann N, McIntosh JR, Howell KE: Direct continuities between cisternae at different levels of the Golgi complex in glucose-stimulated mouse islet beta cells. *PNAS* 2004; 101(15):5565 – 5570.
95. Arvan P, Halban PA: Sorting ourselves out: seeking consensus on trafficking in the beta-cell. *Traffic* 2004;5(1):53 – 61.
96. Morgan DJ, Wei S, Gomes I, et al.: The propeptide precursor proSAAS is involved in fetal neuropeptide processing and body weight regulation. *Journal of Neurochemistry* 2010;113(5): 1275 – 1284.
97. Alarcon C, Verchere CB, Rhodes CJ: Translational control of glucose-induced islet amyloid polypeptide production in pancreatic islets. *Endocrinology* 2012;153(5):2082 – 2087.