International Textbook of Diabetes Mellitus, 4th Ed., Chapter 26 - The genetics of type 2 diabetes

References

1. Tuomi T, Carlsson A, Li H, et al.: Clinical and genetic characteristics of type 2 diabetes with and without GAD antibodies. Diabetes 1999;48(1):150–157.
2. Neel JV: Diabetes mellitus: a “thrifty” genotype rendered detrimental by “progress”? American Journal of Human Genetics 1962;14:353–362.
3. Kaprio J, Tuomilehto J, Koskenvuo M, et al.: Concordance for type 1 (insulin-dependent) and type 2 (non-insulin-dependent) diabetes mellitus in a population-based cohort of twins in Finland. Diabetologia 1992;35(11):1060–1067.
4. Newman B, Selby JV, King MC, et al.: Concordance for type 2 (non-insulin-dependent) diabetes mellitus in male twins. Diabetologia 1987;30(10):763–768.
5. Poulsen P, Kyvik KO, Vaag A, Beck-Nielsen H: Heritability of type II (non-insulin-dependent) diabetes mellitus and abnormal glucose tolerance: a population-based twin study. Diabetologia 1999;42(2):139–145.
6. Medici F, HawaM, Ianari A, et al.:Concordance rate for type II diabetes mellitus in monozygotic twins: actuarial analysis. Diabetologia 1999;42(2):146–150.
7. Meigs JB, Cupples LA, Wilson PW: Parental transmission of type 2 diabetes: the Framingham Offspring Study. Diabetes 2000;49(12):2201–2207.
8. Diamond J: The double puzzle of diabetes. Nature 2003;423(6940): 599–602.
9. Groop L, Forsblom C, Lehtovirta M, et al.:Metabolic consequences of a family history of NIDDM (the Botnia study): evidence for sex-specific parental effects. Diabetes 1996;45(11):1585–1593.
10. Froguel P, Vaxillaire M, Sun F, et al.: Close linkage of glucokinase locus on chromosome 7p to early-onset non-insulin-dependent diabetes mellitus. Nature 1992;356(6365):162–164.
11. YamagataK,Oda N, Kaisaki PJ, et al.: Mutations in the hepatocyte nuclear factor-1alpha gene in maturity-onset diabetes of the young (MODY3). Nature 1996;384(6608):455–458.
12. Yamagata K, Furuta H, Oda N, et al.: Mutations in the hepatocyte nuclear factor-4alpha gene in maturity-onset diabetes of the young (MODY1). Nature 1996;384(6608):458–460.
13. Voight BF, Scott LJ, Steinthorsdottir V, et al.: Twelve type 2 diabetes susceptibility loci identified through large-scale association analysis. Nature Genetics 2010;42(7):579–589.
14. Horikawa Y, Oda N, Cox NJ, et al.: Genetic variation in the gene encoding calpain-10 is associated with type 2 diabetes mellitus. Nature Genetics 2000;26(2):163–175.
15. SongY,Niu T, Manson JE, et al.:Are variants in the CAPN10 gene related to risk of type 2 diabetes? A quantitative assessment of population and family-based association studies. American Journal of Human Genetics 2004;74(2):208–222.
16. Weedon MN, Schwarz PE, Horikawa Y, et al.: Meta-analysis and a large association study confirm a role for calpain-10 variation in type 2 diabetes susceptibility. American Journal of Human Genetics 2003;73(5):1208–1212.
17. Duggirala R, Blangero J, Almasy L, et al.: Linkage of type 2 diabetes mellitus and of age at onset to a genetic location on chromosome 10q in Mexican Americans. American Journal of Human Genetics 1999;64(4):1127–1140.
18. Reynisdottir I,Thorleifsson G, Benediktsson R, et al.: Localization of a susceptibility gene for type 2 diabetes to chromosome 5q34-q35.2. American Journal of Human Genetics 2003;73(2):\ 323–335.

19. Grant SF, Thorleifsson G, Reynisdottir I, et al.: Variant of transcription factor 7-like 2 (TCF7L2) gene confers risk of type 2 diabetes. Nature Genetics 2006;38(3):320–323.
20. Tong Y, Lin Y, Zhang Y, et al.: Association between TCF7L2 gene polymorphisms and susceptibility to type 2 diabetes mellitus: a large Human Genome Epidemiology (HuGE) review and meta-analysis. BMC Medical Genetics 2009;10:15.
21. Reich DE, Lander ES: On the allelic spectrum of human disease. Trends in Genetics 2001;17(9):502–510.
22. Lander ES: The new genomics: global views of biology. Science 1996;274(5287):536–539.
23. Deeb SS, Fajas L, Nemoto M, et al.: A Pro12Ala substitution in PPARgamma2 associated with decreased receptor activity, lower body mass index and improved insulin sensitivity. Nature Genetics 1998;20(3):284–287.
24. Rosengren AH, Braun M, Mahdi T, et al.: Reduced insulin exocytosis in human pancreatic beta-cells with gene variants linked to type 2 diabetes. Diabetes 2012;61(7):1726–1733.
25. Sladek R, Rocheleau G, Rung J, et al.: A genome-wide association study identifies novel risk loci for type 2 diabetes. Nature 2007;445(7130):881–885.
26. Saxena R, Voight BF, Lyssenko V, et al.: Genome-wide association analysis identifies loci for type 2 diabetes and triglyceride levels. Science 2007;316(5829):1331–1336.
27. Scott LJ, Mohlke KL, Bonnycastle LL, et al.: A genome-wide association study of type 2 diabetes in Finns detects multiple susceptibility variants. Science 2007;316(5829):1341–1345.
28. Steinthorsdottir V, Thorleifsson G, Reynisdottir I, et al.: A variant in CDKAL1 influences insulin response and risk of type 2 diabetes. Nature Genetics 2007;39(6):770–775.
29. Zeggini E, Weedon MN, Lindgren CM, et al.: Replication of genome-wide association signals in UK samples reveals risk loci for type 2 diabetes. Science 2007;316(5829):1336–1341.
30. Dina C, Meyre D, Gallina S, et al.: Variation in FTO contributes to childhood obesity and severe adult obesity. Nature Genetics 2007;39(6):724–726.
31. Frayling TM, Timpson NJ, Weedon MN, et al.: A common variant in the FTO gene is associated with body mass index and predisposes to childhood and adult obesity. Science 2007; 316(5826):889–894.
32. Unoki H, Takahashi A, Kawaguchi T, et al.: SNPs in KCNQ1 are associated with susceptibility to type 2 diabetes in East Asian and European populations. Nature Genetics 2008;40(9):1098–1102.
33. Yasuda K, Miyake K, Horikawa Y, et al.: Variants in KCNQ1 are associated with susceptibility to type 2 diabetes mellitus. Nature Genetics 2008;40(9):1092–1097.
34. Kong A, Steinthorsdottir V, Masson G, et al.: Parental origin of sequence variants associated with complex diseases. Nature 2009;462(7275):868–874.
35. DIAbetes Genetics Replication and Meta-analysis (DIAGRAM) Consortium: Genome-wide trans-ancestry meta-analysis provides insight into the genetic architecture of type 2 diabetes susceptibility. Nature Genetics 2014;46(3):234–244.
36. Cho YS, Chen CH, Hu C, et al.: Meta-analysis of genome-wide association studies identifies eight new loci for type 2 diabetes in east Asians. Nature Genetics 2012;44(1):67–72.

37. Imamura M, Maeda S, Yamauchi T, et al.: A single-nucleotide polymorphism in ANK1 is associated with susceptibility to type 2 diabetes in Japanese populations. Human Molecular Genetics 2012;21(13):3042–3049.
38. Kooner JS, Saleheen D, Sim X, et al.: Genome-wide association study in individuals of South Asian ancestry identifies six new type 2 diabetes susceptibility loci. Nature Genetics 2011;43(10):984–989.
39. LiH,GanW, Lu L, et al.: A genome-wide association study identifies GRK5 and RASGRP1 as type 2 diabetes loci in Chinese Hans. Diabetes 2013;62(1):291–298.
40. Palmer ND, McDonough CW, Hicks PJ, et al.: A genome-wide association search for type 2 diabetes genes in African Americans. PLoS One 2012;7(1):e29202.
41. Parra EJ, Below JE, Krithika S, et al.: Genome-wide association study of type 2 diabetes in a sample from Mexico City and a meta-analysis of a Mexican-American sample from Starr County, Texas. Diabetologia 2011;54(8):2038–2046.
42. Shu XO, Long J, CaiQ, et al.: Identification of new genetic risk variants for type 2 diabetes. PLoS Genetics 2010;6(9):e1001127.
43. Sim X, Ong RT, Suo C, et al.: Transferability of type 2 diabetes implicated loci in multi-ethnic cohorts from Southeast Asia. PLoS Genetics 2011;7(4):e1001363.
44. Yamauchi T, Hara K, Maeda S, et al.: A genome-wide association study in the Japanese population identifies susceptibility loci for type 2 diabetes atUBE2E2 and C2CD4A-C2CD4B.Nature Genetic 2010;42(10):864–868.
45. Ma RC, Hu C, TamCH, et al.:Genome-wide association study in a Chinese population identifies a susceptibility locus for type 2 diabetes at 7q32 near PAX4. Diabetologia 2013;56(6):1291–1305.
46. Morris AP,Voight BF, Teslovich TM, et al.: Large-scale association analysis provides insights into the genetic architecture and pathophysiology of type 2 diabetes.Nature Genetics 2012;44(9):981–990.
47. Steinthorsdottir V, Thorleifsson G, Sulem P, et al.: Identification of low-frequency and rare sequence variants associated with elevated or reduced risk of type 2 diabetes. Nature Genetics 2014;46(3):294–298.
48. Lyssenko V, Nagorny CL, Erdos MR, et al.: Common variant in MTNR1B associated with increased risk of type 2 diabetes and impaired early insulin secretion. Nature Genetics 2009;41(1):82–88.
49. Bouatia-Naji N, Bonnefond A, Cavalcanti-Proenca C, et al.: A variant near MTNR1B is associated with increased fasting plasma glucose levels and type 2 diabetes risk. Nature Genetics 2009;41(1):89–94.
50. Prokopenko I, Langenberg C, Florez JC, et al.:Variants in MTNR1B influence fasting glucose levels. Nature Genetics 2009;41(1):77–81.
51. Rung J, Cauchi S, Albrechtsen A, et al.: Genetic variant near IRS1 is associated with type 2 diabetes, insulin resistance and hyperinsulinemia. Nature Genetics 2009;41(10):1110–1115.
52. Saxena R, Hivert MF, Langenberg C, et al.: Genetic variation in GIPR influences the glucose and insulin responses to an oral glucose challenge. Nature Genetics 42(2):142–148.
53. Lyssenko V, Eliasson L, Kotova O, et al.: Pleiotropic effects of GIP on islet function involves osteopontin. Diabetes 2011; 60(9):2424–2433.
54. Ahlqvist E, Osmark P, Kuulasmaa T, et al.: A link between GIP and osteopontin in adipose tissue and insulin resistance. Diabetes 2013;62(6):2088–2094.

55. Ingelsson E, Langenberg C, Hivert MF, et al.: Detailed physiologic characterization reveals diverse mechanisms for novel genetic Loci regulating glucose and insulin metabolism in humans. Diabetes 2010;59(5):1266–1275.
56. Grarup N, Andersen G, Krarup NT, et al.: Association testing of novel type 2 diabetes risk alleles in the JAZF1, CDC123/CAMK1D, TSPAN8, THADA, ADAMTS9, and NOTCH2 loci with insulin release, insulin sensitivity, and obesity in a population-based sample of 4,516 glucose-tolerant middle-aged Danes. Diabetes 2008;57(9):2534–2540.
57. Dimas AS, Lagou V, Barker A, et al.: Impact of type 2 diabetes susceptibility variants on quantitative glycemic traits reveals mechanistic heterogeneity. Diabetes 2013. Epub 2013/12/02.
58. Visscher PM, Hill WG, Wray NR: Heritability in the genomics era: concepts and misconceptions. Nature Reviews Genetics 2008;9(4):255–266.
59. Plomin R, Haworth CM, Davis OS: Common disorders are quantitative traits. Nature Reviews Genetics 2009;10(12):872–878.
60. Mitchell KJ: What is complex about complex disorders? Genome Biology 2012;13(1):237.
61. Yang J, Benyamin B,McEvoy BP, et al.: Common SNPs explain a large proportion of the heritability for human height. NatureGenetics 2010;42(7):565–569.
62. Craddock N, Hurles ME, Cardin N, et al.: Genome-wide association study of CNVs in 16,000 cases of eight common diseases and 3,000 shared controls. Nature 2010;464(7289):713–720.
63. Zeggini E, Scott LJ, Saxena R, et al.: Meta-analysis of genome-wide association data and large-scale replication identifies additional susceptibility loci for type 2 diabetes. Nature Genetics 2008;40(5):638–645.
64. Davidson BL, McCray PB, Jr,: Current prospects for RNA interference-based therapies. Nature Reviews Genetics 2011;12(5): 329–340.
65. Fernandez-Valverde SL, Taft RJ, Mattick JS: MicroRNAs in beta-cell biology, insulin resistance, diabetes and its complications. Diabetes 2011;60(7):1825–1831.
66. Hariharan M, Scaria V, Brahmachari SK: dbSMR: a novel resource of genome-wide SNPs affecting microRNA mediated regulation. BMC Bioinformatics 2009;10:108.
67. Broadbent HM, Peden JF, Lorkowski S, et al.: Susceptibility to coronary artery disease and diabetes is encoded by distinct, tightly linked SNPs in the ANRIL locus on chromosome 9p. Human Molecular Genetics 2008;17(6):806–814.
68. Phillips PC: Epistasis: the essential role of gene interactions in the structure and evolution of genetic systems. Nature Reviews Genetics 2008;9(11):855–867.
69. Cordell HJ, Wedig GC, Jacobs KB, Elston RC: Multilocus linkage tests based on affected relative pairs. American Journal of Human Genetics 2000;66(4):1273–1286.
70. Cox NJ, Frigge M, Nicolae DL, et al.: Loci on chromosomes 2 (NIDDM1) and 15 interact to increase susceptibility to diabetes in Mexican Americans. Nature Genetics 1999;21(2):213–215.
71. Gonzalez-Perez A, Gayan J, Marin J, et al.: Whole-genome conditional two-locus analysis identifies novel candidate genes for late-onset Parkinson's disease. Neurogenetics 2009;10(3):173–181.
72. Nair RP, Duffin KC, Helms C, et al.: Genome-wide scan reveals association of psoriasis with IL-23 and NF-kappaB pathways. Nature Genetics 2009;41(2):199–204.

73. Evans DM, Marchini J, Morris AP, Cardon LR: Two-stage two-locus models in genome-wide association. PLoS Genetics 2006;2(9):e157.
74. Zuk O, Hechter E, Sunyaev SR, Lander ES:The mystery of missing heritability: genetic interactions create phantom heritability. PNAS 2012;109(4):1193–1198.
75. Chong S, Whitelaw E: Epigenetic germline inheritance. Current Opinion in Genetics & Development 2004;14(6):692–696.
76. Liu L, Li Y, Tollefsbol TO: Gene-environment interactions and epigenetic basis of human diseases. Current Issues in Molecular Biology 2008;10(1–2):25–36.
77. Brasacchio D, Okabe J, Tikellis C, et al.: Hyperglycemia induces a dynamic cooperativity of histone methylase and demethylase enzymes associated with gene-activating epigenetic marks that coexist on the lysine tail. Diabetes 2009;58(5):1229–1236.
78. Small KS, Hedman AK, Grundberg E, et al.: Identification of an Imprinted master trans regulator at the KLF14 locus related to multiple metabolic phenotypes. Nature Genetics 2011;43(6):561–564.
79. Bergholdt R, Brorsson C, Palleja A, et al.: Identification of novel type 1 diabetes candidate genes by integrating genome-wide association data, protein-protein interactions, and human pancreatic islet gene expression. Diabetes 2012;61(4):954–962.
80. Suhre K, Shin SY, Petersen AK, et al.: Human metabolic individuality in biomedical and pharmaceutical research. Nature 2011; 477(7362):54–60.
81. Kanehisa M, Goto S, Kawashima S, et al.:The KEGG resource for deciphering the genome. Nucleic Acids Research 2004;32(Database issue):D277–280.
82. Ashburner M, Ball CA, Blake JA, et al.: Gene ontology: tool for the unification of biology. The Gene Ontology Consortium. Nature Genetics 2000;25(1):25–29.
83. Calvano SE, Xiao W, Richards DR, et al.: A network-based analysis of systemic inflammation in humans. Nature 2005;437(7061):1032–1037.
84. Nibbe RK, Koyuturk M, Chance MR: An integrative-omics approach to identify functional sub-networks in human colorectal cancer. PLoS Computational Biology 2010;6(1):e1000639.
85. Pujana MA, Han JD, Starita LM, et al.: Network modeling links breast cancer susceptibility and centrosome dysfunction. Nature Genetics 2007;39(11):1338–1349.
86. Taneera J, Lang S, Sharma A, et al.: A systems genetics approach identifies genes and pathways for type 2 diabetes in human islets. Cell Metabolism 2012;16(1):122–134.
87. Saxena R, Hivert MF, Langenberg C, et al.: Genetic variation in GIPR influences the glucose and insulin responses to an oral glucose challenge. Nature Genetics 2010;42(2):142–148.
88. axena R, Elbers CC, Guo Y, et al.: Large-scale gene-centric meta-analysis across 39 studies identifies type 2 diabetes loci. American Journal of Human Genetics 2012;90(3):410–425.
89. Takeuchi F, Serizawa M, Yamamoto K, et al.: Confirmation of multiple risk Loci and genetic impacts by a genome-wide association study of type 2 diabetes in the Japanese population. Diabetes 2009;58(7):1690–1699.
90. Perry JR, Voight BF, Yengo L, et al.: Stratifying type 2 diabetes cases by BMI identifies genetic risk variants in LAMA1 and enrichment for risk variants in lean compared to obese cases. PLoS Genetics 2012;8(5):e1002741.
91. Huang J, Ellinghaus D, Franke A, et al.: 1000 Genomes-based imputation identifies novel and refined associations for the Wellcome Trust Case Control

Consortium phase 1 Data. European Journal of Human Genetics 2012;20(7):801–805.
92. Timpson NJ, Lindgren CM, Weedon MN, et al.: Adiposity-related heterogeneity in patterns of type 2 diabetes susceptibility observed in genome-wide association data. Diabetes 2009;58(2):505–510.
93. The Wellcome Trust Case Control Consortium: Genome-wide association study of 14,000 cases of seven common diseases and 3,000 shared controls. Nature 2007;447(7145):661–678.
94. Hara K, Fujita H, Johnson TA, et al.: Genome-wide association study identifies three novel loci for type 2 diabetes. Human Molecular Genetics 2014;23(1):239–246.
95. Gudmundsson J, Sulem P, Steinthorsdottir V, et al.: Two variants on chromosome 17 confer prostate cancer risk, and the one in TCF2 protects against type 2 diabetes. Nature Genetics 2007; 39(8):977–983.
96. WincklerW, Burtt NP,Holmkvist J, et al.: Association of common variation in the HNF1 alpha gene region with risk of type 2 diabetes. Diabetes 2005;54(8):2336–2342.
97. AlmindK, Bjorbaek C,Vestergaard H, et al.: Amino acid polymorphisms of insulin receptor substrate-1 in non-insulin-dependent diabetes mellitus. Lancet 1993;342(8875):828–832.
98. Hani EH, Boutin P, Durand E, et al.: Missense mutations in the pancreatic islet beta cell inwardly rectifying K+ channel gene (KIR6.2/BIR): a meta-analysis suggests a role in the polygenic basis of Type II diabetes mellitus in Caucasians. Diabetologia 1998;41(12):1511–1515.
99. Tsai FJ, Yang CF, Chen CC, et al.:A genome-wide association study identifies susceptibility variants for type 2 diabetes inHan Chinese. PLoS Genetics 2010;6(2):e1000847.
100. Dupuis J, Langenberg C, Prokopenko I, et al.: New genetic loci implicated in fasting glucose homeostasis and their impact on type 2 diabetes risk. Nature Genetics 2010;42(2):105–116.
101. Qi L, Cornelis MC, Kraft P, et al.: Genetic variants at 2q24 are associated with susceptibility to type 2 diabetes. Human Molecular Genetics 2010;19(13):2706–2715.
102. Saxena R, Saleheen D, Been LF, et al.: Genome-wide association study identifies a novel locus contributing to type 2 diabetes susceptibility in Sikhs of Punjabi origin from India. Diabetes 2013; 62(5):1746–1755.
103. Salonen JT, Uimari P, Aalto JM, et al.: Type 2 diabetes whole-genome association study in four populations: the DiaGen consortium. American Journal of Human Genetics 2007;81(2): 338–345.
104. Tabassum R, Chauhan G, Dwivedi OP, et al.: Genome-wide association study for type 2 diabetes in Indians identifies a new susceptibility locus at 2q21. Diabetes 2013;62(3):977–986.
105. Minton JA, Hattersley AT, Owen K, et al.: Association studies of genetic variation in the WFS1 gene and type 2 diabetes in U.K. populations. Diabetes 2002;51(4):1287–1290.
106. Sandhu MS, Weedon MN, Fawcett KA, et al.: Common variants in WFS1 confer risk of type 2 diabetes. Nature Genetics 2007;39(8):951–953.