

International Textbook of Diabetes Mellitus, 4th Ed., Chapter 30 - Molecular genetics of type 1 diabetes

References

1. Diabetes Epidemiology Research International Group: Geographic patterns of childhood insulin-dependent diabetes mellitus. *Diabetes* 1988;37:1113–1119.
2. Davies JL, Kawaguchi Y, Bennett ST, et al.: A genome-wide search for human type 1 diabetes susceptibility genes. *Nature* 1994;371: 130–136.
3. Steck AK, RewersMJ: Genetics of type 1 diabetes. *Clinical Chemistry* 2011;57: 176–185.
4. Redondo MJ, Jeffrey J, Fain PR, et al.: Concordance for islet autoimmunity among monozygotic twins. *New England Journal of Medicine* 2008;359:2849–2850.
5. Patterson CC, Dahlquist GG, Gyurus E, et al., EURODIAB Study Group: Incidence trends for childhood type 1 diabetes in Europe during 1989–2003 and predicted new cases 2005–20: a multicentre prospective registration study. *Lancet* 2009;373:2027–2033.
6. Onkamo P, Vaananen S, Karvonen M, Tuomilehto J: Worldwide increase in incidence of type 1 diabetes: the analysis of data on published incidence trends. *Diabetologia* 1999;42:1395–1403.
7. Feltbower RG, Bodansky HJ, McKinney PA, et al.: Trends in the incidence of childhood diabetes in south Asians and other children in Bradford, UK. *Diabetic Medicine* 2002;19:162–166.
8. Ji J, Hemminki K, Sundquist J, Sundquist K: Ethnic differences in incidence of type 1 diabetes among second-generation immigrants and adoptees from abroad. *Journal of Clinical Endocrinology & Metabolism* 2010;95:847–850.
9. Harron KL, McKinney PA, Feltbower RG, et al.: Incidence rate trends in childhood type 1 diabetes in Yorkshire, UK 1978–2007: effects of deprivation and age at diagnosis in the south Asian and non south Asian populations. *Diabetic Medicine* 2011;28:1508–1513.
10. Ekeh S, Popovic P, Muntoni S, et al.: Incidence of diabetes mellitus among children of Italian migrants substantiates the role of genetic factors in the pathogenesis of type 1 diabetes. *European Journal of Pediatrics* 2009;168: 613–617.
11. Barrett JC, Clayton D, Concannon P, et al.: Genome-wide association study and meta-analysis find over 40 loci affect risk of type 1 diabetes. *Nature Genetics* 2009;41:703–707.
12. Singal DP, Blajchman MA: Histocompatibility (HL-A) antigens, lymphocytotoxic antibodies and tissue antibodies in patients with diabetes mellitus. *Diabetes* 1973;22:429–432.
13. Rotter JI, Landaw EM: Measuring the genetic contribution of a single locus to a multilocus disease. *Clinical Genetics* 1984;26:529–542.
14. Noble JA, Valdes AM: Genetics of the HLA region in the prediction of type 1 diabetes. *Current Diabetes Reports* 2011;11:533–542.
15. Rewers M, Bugawan TL, Norris JM, et al.: Newborn screening for HLA markers associated with IDDM: Diabetes Autoimmunity Study in the Young (DAISY). *Diabetologia* 1996;39:807–812.
16. Thomson G, Valdes AM, Noble JA, et al.: Relative predispositional effects of HLA class II DRB1-DQB1 haplotypes and genotypes on type 1 diabetes: a meta-analysis. *Tissue Antigens* 2007;70:110–127.

17. Roep BO, Schipper R, Verduyn W, et al.: HLA-DRB1*0403 is associated with dominant protection against IDDM in the general Dutch population and subjects with high-risk DQA1*0301-DQB1*0302/DQA1*0501-DQB1*0201 genotype. *Tissue Antigens* 1999;54:88–90.
18. Baisch JM, Weeks T, Giles R, et al.: Analysis of HLA-DQ genotypes and susceptibility in insulin-dependent diabetes mellitus. *New England Journal of Medicine* 1990;322:1836–1841.
19. Redondo MJ, Kawasaki E, Mulgrew CL, et al.: DR- and DQ-associated protection from type 1A diabetes: comparison of DRB1*1401 and DQA1*0102-DQB1*0602. *Journal of Clinical Endocrinology & Metabolism* 2000;85:3793–3797.
20. Pugliese A, Gianani R, Moromisato R, et al.: HLA-DQB1*0602 is associated with dominant protection from diabetes even among islet cell antibody-positive first-degree relatives of patients with IDDM. *Diabetes* 1995;44:608–613.
21. Pugliese A, Kawasaki E, Zeller M, et al.: Sequence analysis of the diabetes protective human leukocyte antigen-DQB1*0602 allele in unaffected, islet cell antibody-positive first-degree relatives and in rare patients with type 1 diabetes. *Journal of Clinical Endocrinology & Metabolism* 1999;84:1722–1728.
22. Rayner ML, Kelly MA, Mijovic CH, Barnett AH: Sequencing of the second exon of the MHC class II DQ6 alleles in patients with type 1 diabetes. *Autoimmunity* 2002;35:155–157.
23. Valdes AM, Thomson G, Graham J, et al.: D6S265*15 marks a DRB1*15,DQB1*0602 haplotype associated with attenuated protection from type 1 diabetes mellitus. *Diabetologia* 2005; 48:2540–2543.
24. Husain Z, Kelly MA, Eisenbarth GS, et al.: The MHC type 1 diabetes susceptibility gene is centromeric to HLA-DQB1. *Journal of Autoimmunity* 2008;10:596–600.
25. Price P, Cheong KYM, Boodhoo A, et al.: Can MHC class II genes mediate resistance to type 1 diabetes? *Immunology and Cell Biology* 2001;79:602–606.
26. Abraham RS, Wen L, Marietta EV, David CS: Type 1 diabetes predisposing MHC alleles influence the selection of glutamic acid decarboxylase (GAD) 65-specific T cells in a transgenic model. *Journal of Immunology* 2001;166:1370–1379.
27. Abraham RS, Kudva YC, Wilson SB, et al.: Co-expression of HLA DR3 and DQ8 results in the development of spontaneous insulinitis and loss of tolerance to GAD65 in transgenic mice. *Diabetes* 2000;49:548–554.
28. Wen L, Wong FS, Tang J, et al.: In vivo evidence for the contribution of human histocompatibility leukocyte antigen (HLA)-DQ molecules to the development of diabetes. *Journal of Experimental Medicine* 2000;191:97–104.
29. Wen L, Chen N-Y, Tang J, et al.: The regulatory role of DR4 in a spontaneous diabetes DQ8 transgenic model. *Journal of Clinical Investigation* 2001;107:871–880.
30. Rajagopalan G, Kudva YC, Chen L, et al.: Autoimmune diabetes in HLA-DR3/DQ8 transgenic mice expressing the co-stimulatory molecule B7-1 in the beta cells of islets of Langerhans. *International Immunology* 2003;15:1035–1044.
31. Kudva YC, Rajagopalan G, Raju R, et al.: Modulation of insulinitis and type 1 diabetes by transgenic HLA-DR3 and DQ8 in NOD mice lacking endogenous MHC class II. *Human Immunology* 2002;63:987–999.
32. Cucca F, Lampis R, Congia M, et al.: A correlation between the relative predisposition of MHC class II alleles to type 1 diabetes and the structure of their proteins. *Human Molecular Genetics* 2001;10:2025–2037.
33. Wherrett DK, Singer SM, McDevitt HO: Reduction in diabetes incidence in an I-Ag7 transgenic nonobese diabetic mouse line. *Diabetes* 1997;46:1970–1974.

34. Varney MD, Valdes AM, Carlson JA, et al.: HLA DPA1,DPB1 alleles and haplotypes contribute to the risk associated with type 1 diabetes: analysis of the Type 1 Diabetes Genetics Consortium families. *Diabetes* 2010;59:2055–2062.
35. Noble JA, Valdes AM, Varney MD, et al.: HLA class I and genetic susceptibility to type 1 diabetes: results from the Type 1 Diabetes Genetics Consortium. *Diabetes* 2010;59:2972–2979.
36. Howson JMM, Walker NM, Clayton D, Todd JA; the Type 1 Diabetes Genetics Consortium: Confirmation of HLA class II independent type 1 diabetes associations in the major histocompatibility complex including HLA-B and HLA-A. *Diabetes, Obesity and Metabolism* 2009;11:31–45.
37. Fourlanos S, Varney MD, Tait BD, et al.: The rising incidence of type 1 diabetes is accounted for by cases with lower-risk human leukocyte antigen genotypes. *Diabetes Care* 2008;31:1546–1549.
38. Barratt BJ, Payne F, Lowe CE, et al.: Remapping the insulin gene/IDDM2 locus in type 1 diabetes. *Diabetes* 2004;53:1884–1889.
39. Vafiadis P, Bennett ST, Todd JA, et al.: Insulin expression in human thymus is modulated by INS VNTR alleles at the IDDM2 locus. *Nature Genetics* 1997;15:289–292.
40. Vafiadis P, Ounissi-Benkhalha H, Palumbo M, et al.: Class III alleles of the variable number of tandem repeat insulin polymorphism associated with silencing of thymic insulin predispose to type 1 diabetes. *Journal of Clinical Endocrinology & Metabolism* 2001;86:3705–3710.
41. Ueda H, Howson JM, Esposito L, et al.: Association of the T-cell regulatory gene CTLA4 with susceptibility to autoimmune disease. *Nature* 2003;423:506–511.
42. Anjos SM, Shao W, Marchand L, Polychronakos C: Allelic effects on gene regulation at the autoimmunity-predisposing CTLA4 locus: a re-evaluation of the 3' +6230G>A polymorphism. *Genes & Immunity* 2005;6:6305–6311.
43. QuHq, Bradfield JP, Grant SF, et al.; Type 1 Diabetes Genetics Consortium: Remapping the type 1 diabetes association of the CTLA4 locus. *Genes & Immunity* 2009;10(Suppl 1):S27–S32.
44. Steck AK, Liu SY, McFann K, et al.: Association of the PTPN22/LYP gene with type 1 diabetes. *Pediatric Diabetes* 2006;7:274–278.
45. Vang T, Congia M, Macis MD, et al.: Autoimmune-associated lymphoid tyrosine phosphatase is a gain-of-function variant. *Nature Genetics* 2005;37:1317–1319.
46. Vella A, Cooper JD, Lowe CE, et al.: Localization of a type 1 diabetes locus in the IL2RA/CD25 region by use of tag single-nucleotide polymorphisms. *American Journal of Human Genetics* 2005;76:773–779.
47. Lowe CE, Cooper JD, Brusko T, et al.: Large-scale genetic fine-mapping and genotype-phenotype associations implicate polymorphism in the IL2RA region in type 1 diabetes. *Nature Genetics* 2007;39:1074–1082.
48. Dendrou CA, Plagnol V, Fung E, et al.: Cell-specific protein phenotypes for the autoimmune locus IL2RA using a genotype-selectable human bioresource. *Nature Genetics* 2009;41:1011–1015.
49. Garg G, Tyler JR, Yang JHM, et al.: Type 1 diabetes-associated IL2RA variation lowers IL 2 signaling and contributes to diminished CD4+ CD25+ regulatory T cell function. *Journal of Immunology* 2012;188:4644–4653.
50. Concannon P, Chen W-M, Julier C, et al.: Genome-wide scan for linkage to type 1 diabetes in 2,496 multiplex families from the Type 1 Diabetes Genetics Consortium. *Diabetes* 2009;58:1018–1022.

51. Smyth DJ, Cooper JD, Bailey R, et al.: A genome-wide association study of nonsynonymous SNPs identifies a type 1 diabetes locus in the interferon-induced helicase (IFIH1) region. *Nature Genetics* 2006;38:617–619.
52. Wellcome Trust Case Control Consortium: Genome-wide association study of 14,000 cases of seven common diseases and 3,000 shared controls. *Nature* 2007;447:661–678.
53. Bradfield JP, Qu H-Q, Wang K, et al.: A genome-wide meta-analysis of six type 1 diabetes cohorts identifies multiple associated loci. *PLoS Genetics* 2011;7:e1002293.
54. Guo D, Li M, Zhang Y, et al.: A functional variant of SUMO4, a new I kappa B alpha modifier, is associated with type 1 diabetes. *Nature Genetics* 2004;36:837–841.
55. Baniasadi V, Das SN: No evidence for association of PTPN22 R620W functional variant C1858T with type 1 diabetes in Asian Indians. *Journal of Cellular and Molecular Medicine* 2008;12:1061–1062.
56. Nejentsev S, Walker N, Riches D, et al.: Rare variants of IFIH1, a gene implicated in antiviral response, protect against type 1 diabetes. *Science* 2009;324:387–389.
57. Chistiakov DA, Voranova NV, Savost'Anov KV, Turakulov RI: Loss-of-function mutations E627X and I923V of IFIH1 are associated with lower poly(I:C)-induced interferon- β production in peripheral blood mononuclear cells of type 1 diabetes patients. *Human Immunology* 2010;71:1128–1134.
58. Clayton DG: Prediction and interaction in complex disease genetics: experience in type 1 diabetes. *PLoS Genetics* 2009;5:e1000540.
59. Bonifacio E, Hummel M, Walter M, et al.: IDDM1 and multiple family history of type 1 diabetes combine to identify neonates at high risk for type 1 diabetes. *Diabetes Care* 2004;27:2695–2700.
60. TEDDY Study Group: The Environmental Determinants of Diabetes in the Young (TEDDY) study: study design. *Pediatric Diabetes* 2007;8:286–98.
61. Aly TA, Ide A, Jahromi MM, et al.: Extreme genetic risk for type 1A diabetes. *PNAS* 2006;103:14074–14079.
62. Vidovic D, Xie Y, Rinderspacher A, et al.: Distinct functional and conformational states of the human lymphoid tyrosine phosphatase catalytic domain can be targeted by choice of the inhibitor chemotype. *Journal of Computer-Aided Molecular Design* 2011; 25:873–883.
63. Yang XY, Levi D, Ounissi-Benkalha H, et al.: Screening for novel lead compounds increasing insulin expression in medullary thymic epithelial cells. *European Journal of Pharmacology* 2012;688:84–89.
64. Erlich HA, Valdes AM, Noble J, et al.: HLADR-DQ haplotypes and genotypes and type 1 diabetes risk: analysis of the Type 1 Diabetes Genetics Consortium families. *Diabetes* 2008;57:1084–1092.
65. Sugihara S, Ogata T, Kawamura T, et al.: HLA-class II and class I genotypes among Japanese children with type 1A diabetes and their families. *Pediatric Diabetes* 2012;13:33–44.
66. Pociot F, Akolkar B, Concannon P, et al.: Genetics of type 1 diabetes: what's next? *Diabetes* 2010;59:1561–1571.